

Appendix F

Answers to Selected Exercises

1 · Functions and Their Graphs

1.1 · Linear Models

1.1.7 · Linear Models (Homework 1.1)


1.1.7.1.

Answer.

h	0	3	6	9	10
T	65	80	95	110	115

a $T = 65 + 5h$

b



c 95°


d 3 p.m.

1.1.7.3.

Answer.

w	0	4	8	12	16
A	250	190	130	70	10

a $A = 250 - 15w$



c 75 gallons


d Until the fifth week

1.1.7.5.

Answer.

a $P = -800 + 40t$

b $(0, -800), (20, 0)$



c The P -intercept, -800 , is the initial ($t = 0$) value of the profit. Phil and Ernie start out \$800 in debt. The t -intercept, 20 , is the number of hours required for Phil and Ernie to break even.


1.1.7.7.

Answer.

a $C = 5000 + 0.125d$

b Complete the table of values.

Miles Driven	4000	8000	12,000	16,000	20,000
Cost (\$)	5500	6000	6500	7000	7500




d \$500

e More than 16,000 miles


1.1.7.9.

Answer.



1.1.7.11.


Answer.



1.1.7.13.

Answer.


a (8, 0), (0, 4)



1.1.7.15.

Answer.


a (4, 0), (0, -3)



1.1.7.17.

Answer.


a (9, 0), (0, -4)



1.1.7.19.

Answer.


a (-2250, 0), (0, 1500)



1.1.7.21.

Answer.


a (12, 0), (0, 4)



1.1.7.23.

Answer.

a $\left(\frac{3}{2}, 0\right), \left(0, \frac{11}{3}\right)$




1.1.7.25.

Answer.

a $\$2.40x, \$3.20y$

b $2.40x + 3.20y = 19,200$




d The y -intercept, 6000 gallons, is the amount of premium that the gas station owner can buy if he buys no regular. The x -intercept, 8000 gallons, is the amount of regular he can buy if he buys no premium.

1.1.7.27.

Answer.

a $9x$ mg, $4y$ mg

b $9x + 4y = 1800$



d The x -intercept, 200 grams, tells how much fig Delbert should eat if he has no bananas, and the y -intercept, 450 grams, tells how much banana he should eat if he has no figs.

1.1.7.29.

Answer.

- a $(3, 0), (0, 5)$
- b $\left(\frac{1}{2}, 0\right), \left(0, \frac{-1}{4}\right)$
- c $\left(\frac{5}{2}, 0\right), \left(0, \frac{-3}{2}\right)$

1.1.7.31.

Answer.

- a $(0, b)$
- b $\left(\frac{-b}{m}, 0\right)$, if $m \neq 0$


1.1.7.33.

Answer. $-2x + 3y = 2400$

1.1.7.37.

Answer.


a $y = 6 - 2x$



1.1.7.41.

Answer.


a $y = 0.02 - 0.04x$



1.1.7.45.

Answer.

- a $(100, 0), (0, 25)$
- b $y = 25 - \frac{1}{4}x$



d $(p, 0), (0, q)$

e The value of a is the x -intercept, and the value of b is the y -intercept.


1.1.7.35.

Answer. $3x + 400y = 240$

1.1.7.39.

Answer.


a $y = \frac{3}{4}x - 300$



1.1.7.43.

Answer.


a $y = 210 - 35x$



1.1.7.47.

Answer.


- a $(0.04, 0), (0, -0.05)$
- b $y = 1.25x - 0.05$



1.1.7.49.**Answer.**


a $(-60, 0), (0, 12)$

b $y = 12 + \frac{1}{5}x$

**1.1.7.51.****Answer.**

a $(-42, 0), (0, -28)$

b $y = -\frac{2}{3}x - 28$



1.2 · Functions

1.2.10 · Functions (Homework 1.2)

1.2.10.1.**Answer.** Function; the tax is determined by the price of the item.**1.2.10.3.****Answer.** Not a function; incomes may differ for same number of years of education.**1.2.10.5.****Answer.** Function; weight is determined by volume.**1.2.10.7.****Answer.** Input: items purchased; output: price of item. Yes, a function because each item has only one price.**1.2.10.9.****Answer.** Input: topics; output: page or pages on which topic occurs. No, not a function because the same topic may appear in more than one page.**1.2.10.11.****Answer.** Input: students' names; output: students' scores on quizzes, tests, etc. No, not a function because the same student can have different grades on different tests.**1.2.10.13.****Answer.** Input: person stepping on scales; output: person's weight. Yes, a function because a person cannot have two different weights at the same time.**1.2.10.15.****Answer.** No**1.2.10.17.****Answer.** Yes**1.2.10.19.****Answer.** Yes**1.2.10.21.****Answer.** Yes**1.2.10.23.****Answer.** No**1.2.10.25.****Answer.** Yes**1.2.10.27.****Answer.**

a 60

b 37.5

c 30

1.2.10.29.**Answer.**

- a 15%
- b 14%
- c \$7010–\$9169

1.2.10.31.**Answer.**

- a 67.7: In 1985, 67.7% of 20–24 year old women had not yet had children.
- b 1987: Approximately 68% of 20–24 year old women had not yet had children in 1987.
- c $f(1997) = 64.9$

1.2.10.33.**Answer.**

- a No
- b 60; no
- c Decreasing

1.2.10.35.**Answer.**

- a 1991
- b 1 yr
- c 1 yr
- d About 7300

1.2.10.37.**Answer.**

- a Approximately \$1920
- b \$5 or \$15
- c $7.50 < d < 12.50$

1.2.10.39.**Answer.**

- a 1968, about \$8.70
- b 1989, about \$5.10
- c 1967, approximately 1970

1.2.10.41.**Answer.**

- a 0
- b 10
- c -19.4
- d $\frac{14}{3}$

1.2.10.43.**Answer.**

- a 1
- b 6
- c $\frac{3}{8}$
- d 96.48

1.2.10.45.**Answer.**

a $\frac{5}{6}$

b 9

c $\frac{-1}{10}$

d $\frac{12}{13} \approx 0.923$

1.2.10.47.**Answer.**

a $\sqrt{12}$

b 0

c $\sqrt{3}$

d $\sqrt{0.2} \approx 0.447$

1.2.10.49.**Answer.**a $V(12) = 1120$: After 12 years, the SUV is worth \$1120.b $t = 12.5$: The SUV has zero value after $12\frac{1}{2}$ years.

c The value 2 years later

1.2.10.51.**Answer.**a $N(6000) = 2000$: 2000 cars will be sold at a price of \$6000.b $N(p)$ decreases with increasing p because fewer cars will be sold when the price increases.c $2N(D)$ represents twice the number of cars that can be sold at the current price.**1.2.10.53.****Answer.**a $v(250) = 54.8$ is the speed of a car that left 250-foot skid marks.

b $833\frac{1}{3}$ feet

c $v\left(833\frac{1}{3}\right) = 100$

1.2.10.55.**Answer.**

a June 21–24, June 29–July 3, July 8–14

b June 17–21, June 25–29, July 4–7

c June 22–24, June 27, June 29–July 4, July 8–14

d Avalanches occur when temperatures rise above freezing immediately after snowfall.

1.2.10.57.**Answer.**

a No

b Yes

c Moving downwards on the graph corresponds to moving downwards in the ocean.

1.2.10.59.**Answer.**

$$\begin{array}{ll} \text{a } 27a^2 - 18a & \text{c } 3a^2 - 6a + 2 \\ \text{b } 3a^2 + 6a & \text{d } 3a^2 + 6a \end{array}$$

1.2.10.63.**Answer.**

$$\begin{array}{ll} \text{a } 8x^3 - 1 & \text{c } x^6 - 1 \\ \text{b } 2x^3 - 2 & \text{d } x^6 - 2x^3 + 1 \end{array}$$

1.2.10.65.**Answer.**

$$\begin{array}{ll} \text{a } a^6 & \text{c } a^3b^3 \\ \text{b } a^{12} & \text{d } a^3 + 3a^2b + 3ab^2 + b^3 \end{array}$$

1.2.10.61.**Answer.**

$$\begin{array}{ll} \text{a } 8 & \text{c } 8 \\ \text{b } 8 & \text{d } 8 \end{array}$$

1.2.10.67.**Answer.**

$$\begin{array}{llll} \text{a } 96a^5 & \text{c } 3a^{10} \\ \text{b } 6a^5 & \text{d } 9a^{10} \end{array}$$

1.2.10.69.**Answer.**

$$\begin{array}{ll} \text{a } 11 & 4 \\ \text{b } 13 & \text{d } 3a + 3b - 2 \\ \text{c } 3a + 3b - 2 & \end{array}$$

1.2.10.71.**Answer.**

$$\begin{array}{ll} \text{a } 19 & 6 \\ \text{b } 28 & \text{d } a^2 + 2ab + b^2 + 3 \\ \text{c } a^2 + b^2 + 3 & \end{array}$$

1.2.10.73.**Answer.**

$$\begin{array}{ll} \text{a } \frac{\sqrt{3}+1}{2} & \text{c } \frac{\sqrt{a+1}+\sqrt{b+1}}{2} \\ \text{b } \sqrt{6} & \text{d } \sqrt{a+b+1} \end{array}$$

This function does NOT satisfy $f(a+b) = f(a) + f(b)$.

This function does NOT satisfy $f(a+b) = f(a) + f(b)$.

This function does NOT satisfy $f(a+b) = f(a) + f(b)$.

1.2.10.75.**Answer.**

$$\begin{array}{ll} \text{a } \frac{-5}{3} & \frac{-2}{b} \\ \text{b } \frac{-2}{5} & \\ \text{c } \frac{-2}{a} & \text{d } \frac{-2}{a+b} \end{array}$$

This function does NOT satisfy $f(a+b) = f(a) + f(b)$.

1.2.10.77.**Answer.**

a	x	0	10	20	30	40	50	60	70	80	90	100
	$f(x)$	800	840	840	800	720	600	440	240	0	-280	-600

$$a = 50 \text{ and } b = 60$$

b	x	50	51	52	53	54	55	56	57	58
	$f(x)$	600	585.8	571.2	556.2	540.8	525	508.8	492.2	475.2

$$c = 56 \text{ and } d = 57$$

c	x	56	56.1	56.2	56.3	56.4	56.5	56.6
	$f(x)$	508.8	507.158	505.512	503.862	502.208	500.55	498.888

$$p = 56.5 \text{ and } q = 56.6$$

$$d \ s = 56.55$$

$$e \ f(56.55) = 499.7195$$

1.2.10.79.

Answer. 94.85

1.3 · Graphs of Functions

1.3.6 · Graphs of Functions (Homework 1.3)

1.3.6.1.

Answer.

a $-2, 0, 5$

b 2

c $h(-2) = 0, h(1) = 0, h(0) = -2$

d 5

e 3

f Increasing: $(-3, 0)$ and $(1, 3)$; decreasing: $(0, 1)$ and $(3, 5)$

1.3.6.3.

Answer.

a $-1, 2$

b $3, -1.3$

c $R(-2) = 0, R(2) = 0, R(4) = 0, R(0) = 4$

d Max: 4 ; min: -5

e Max at $p = 0$; min at $p = 5$

f Increasing: $(-3, 0)$ and $(1, 3)$; decreasing: $(0, 1)$ and $(3, 5)$

1.3.6.5.

Answer.

a $0, \frac{1}{2}, 0$

b 0.9

c $\frac{-5}{6}, \frac{-1}{6}, \frac{7}{6}, \frac{11}{6}$

d Max: 1 ; min: -1




e Max at $x = -1.5, 0.5$; min at $x = -0.5, 1.5$

1.3.6.7.**Answer.**

a 2, 2, 1

b $-6 \leq s < -4$ or $0 \leq s < 2$

c Max: 2; min: -1

d Max for $-3 \leq s < -1$ or $3 \leq s < 5$; min for $-6 \leq s < -4$ or $0 \leq s < 2$ **1.3.6.9.****Answer.** (a) and (d)**1.3.6.11.****Answer.****1.3.6.13.****Answer.****1.3.6.15.****Answer.****1.3.6.17.****Answer.**

- a $f(1000) = 1495$: The speed of sound at a depth of 1000 meters is approximately 1495 meters per second.
- b $d = 570$ or $d = 1070$: The speed of sound is 1500 meters per second at both a depth of 570 meters and a depth of 1070 meters.
- c The slowest speed occurs at a depth of about 810 meters and the speed is about 1487 meters per second, so $f(810) = 1487$.
- d f increases from about 1533 to 1541 in the first 110 meters of depth, then drops to about 1487 at 810 meters, then rises again, passing 1553 at a depth of about 1600 meters.

1.3.6.19.**Answer.**

- a $f(1985) = 41$: The federal debt in 1985 was about 41% of the gross domestic product.
- b $t = 1942$ or $t = 1955$: The federal debt was 70% of the gross domestic product in 1942 and 1955.
- c In about 1997, the debt was about 67% of the gross domestic product, so $f(1997) \approx 67.3$.
- d The percentage basically dropped from 1946 to 1973, but there were small rises around 1950, 1954, 1958, and 1968, so the longest time interval was from 1958 to 1967.

1.3.6.21.**Answer.**

- a i $x = -3$
 ii $x < -3$
 iii $x > -3$
- b I $x = -3$
 II $x < -3$
 III $x > -3$
- c On the graph of $y = -2x + 6$, a value of y is the same as a value of $-2x + 6$, so parts (a) and (b) are asking for the same x 's.

1.3.6.23.**Answer.**

- a $x = 0.6$
 b $x = -0.4$
 c $x > 0.6$
 d $x < -0.4$

1.3.6.25.**Answer.**

- a $x = -1$ or $x = 1$
 b Approximately $-3 \leq x \leq -2$ or $2 \leq x \leq 3$

1.3.6.27.**Answer.**

- a 3.5
 b $-2.2, -1.2, 3.4$
 c $p < -3.1$ or $0.3 < p < 2.8$
 d $0.5 < B < 5.5$
 e $p < -1.7$ or $p > 1.7$

1.3.6.29.**Answer.**

- a i $-2, 2$
 ii $-2.8, 0, 2.8$
 iii $-2.5 < q < -1.25$ or $1.25 < q < 2.5$
- b $-2 < q < 0$ or $q > 2$

1.3.6.31.**Answer.**

- a He has an error: $f(-3)$ cannot have both the value 5 and also the value -2 , and $f(-1)$ cannot have both values 2 and -4 .
- b Her readings are possible for a function: each input has only one output.

1.3.6.33.**Answer.**


a

x	-4	-2	3	5
$g(x)$	4.5	5.7	5.2	3.3


- b $-4.8, 4.8$

1.3.6.35.**Answer.**

- a $(-1.6, 4.352), (1.6, -4.352)$
- b $F(-1.6) = 4.352; F(1.6) = -4.352$

1.3.6.37.**Answer.**

The curve cannot be distinguished from the x -axis in the standard window because the values of y are closer to zero than the resolution of the calculator can display. The second window provides sufficient resolution to see the curve.

1.3.6.39.**Answer.**

The curve looks like two vertical lines in the standard window because that window covers too small a region of the plane. The second window allows us to see the turning points of the curve.

1.3.6.41.**Answer.**

- a $x = 4$ b $x = -5$ c $x > 1$ d $x < 14$

1.3.6.43.**Answer.**

a $x = 11$

b $x = -10$

c $x \geq -5$

d $x \leq 8$

1.3.6.45.**Answer.**

a $x = 4$

b $x < 22$

1.3.6.47.**Answer.**

a $x = 20$

b $x \leq 7$


1.3.6.49.**Answer.**

a $-15, 5, 20$


b $-13, 2, 22$

1.4 · Slope and Rate of Change

1.4.9 · Slope and Rate of Change (Homework 1.4)

1.4.9.1.**Answer.** Anthony**1.4.9.3.****Answer.** Bob's driveway**1.4.9.5.****Answer.** -1 **1.4.9.7.****Answer.** $-\frac{2}{3}$ **1.4.9.9.****Answer.**


b $\frac{3}{4}$

1.4.9.11.**Answer.**

b -3

1.4.9.13.

Answer.






a

b $\frac{8}{5}$

1.4.9.15.

Answer.



1.4.9.17.

Answer.

- | | | | | |
|---|------|------|--------------------|------------------|
| a | i -3 | ii 6 | iii $-\frac{3}{2}$ | iv $\frac{9}{2}$ |
| b | i -4 | ii 8 | iii $\frac{8}{3}$ | iv $\frac{4}{3}$ |

1.4.9.19.

Answer. $\frac{100}{7}$ ft \approx 14.286 ft \approx 14 ft 3.4 in

1.4.9.21.

Answer.

- | | | | |
|------|-------|------|-----|
| a IV | b III | c II | d I |
|------|-------|------|-----|

1.4.9.23.Answer. $\frac{3}{4}$ **1.4.9.25.**Answer. -4000 **1.4.9.27.**

Answer.

1.4.9.29.

Answer.

a $\frac{5}{2}$

x	y
3	$\frac{7}{2}$
6	11

 -3

x	y
-1	30
5	12


1.4.9.31.

Answer.

t	4	8	20	40
S	32	64	160	320

c 8 dollars/hour

d The typist is paid \$8 per hour.

**1.4.9.33.**

Answer.

a 1250 barrels/day

b The slope indicates that oil is pumped at a rate of 1250 barrels per day.

1.4.9.35.

Answer.

a -6 liters/day

b The slope indicates that the water is diminishing at a rate of 6 liters per day.

1.4.9.37.

Answer.

a 12 inches/foot

b The slope gives the conversion rate of 12 inches per foot.

1.4.9.39.

Answer.

a 4 dollars/kilogram

b The slope gives the unit price of \$4 per kilogram

1.4.9.41.

Answer. (a)

1.4.9.43.

Answer.

a Yes, the slope between any two points is $\frac{1}{2}$.

b 0.5 grams of salt per degree Celsius

1.4.9.45.

Answer.

a Yes

b 2π

1.4.9.47.

Answer.

a $\frac{1500 \text{ meters}}{1 \text{ second}}$

b 3375 meters

1.4.9.49.

Answer.

a The distances are known.

b 5.7 km per second

1.4.9.51.

Answer.

a About 18°C


b 0.3 km to 0.4 km

c About -28°C per kilometer


1.4.9.53.

Answer.

a -3




b 2




1.4.9.55.

Answer.

a $-\frac{1}{4}$



b -1



1.4.9.57.**Answer.**

a $(1, F(1)), (4, F(4)); \quad F(4) - F(1)$

b $(r, f(r)), (s, f(s)); \quad f(s) - f(r)$

1.4.9.59.**Answer.**

a $(2, H(2)), (3, H(3)); \quad H(3) - H(2)$

b $(a, g(a)), (b, g(b)); \quad g(b) - g(a)$

1.4.9.61.**Answer.**

a $(c, s(c)), (d, s(d)); \quad s(c)(d - c)$

b $(x_1, q(x_1)), (x_2, q(x_2)); \quad q(x_2)(x_2 - x_1)$

1.4.9.63.**Answer.**

a $(1, f(1)), (5, f(5)); \quad \frac{f(5) - f(1)}{4}$

b $(-1, f(-1)), (2, f(2)); \quad \frac{f(2) - f(-1)}{3}$

1.4.9.65.**Answer.**

a $(a, f(a)), (b, f(b)); \quad \frac{f(b) - f(a)}{b - a}$

b $(a, f(a)), (a + \Delta x, f(a + \Delta x)); \quad \frac{f(a + \Delta x) - f(a)}{\Delta x}$

1.5 · Linear Functions**1.5.6 · Linear Functions (Homework 1.5)****1.5.6.1.****Answer.**

a $y = \frac{1}{2} - \frac{3}{2}x$

b Slope $\frac{-3}{2}$, y -intercept $\frac{1}{2}$

1.5.6.5.**Answer.**

a $y = -22 + 14x$

b Slope 14, y -intercept -22

1.5.6.3.**Answer.**

a $y = \frac{1}{9} - \frac{1}{6}x$

b Slope $\frac{-1}{6}$, y -intercept $\frac{1}{9}$

1.5.6.7.**Answer.**


a $y = -29$

b Slope 0, y -intercept -29

1.5.6.9.**Answer.**


a $y = \frac{49}{3} - \frac{5}{3}x$

b Slope $-\frac{5}{3}$, y -intercept $\frac{49}{3}$

1.5.6.11.**Answer.**

b $y = -2 + 3x$

c $\frac{2}{3}$

1.5.6.13.**Answer.**

b $y = -6 + \frac{5}{3}x$

c $\frac{-18}{5}$

1.5.6.15.**Answer.** 5**1.5.6.17.****Answer.** $-\frac{1}{4}$ **1.5.6.19.****Answer.** $m = \frac{-A}{B}$, x -intercept $\left(\frac{C}{A}, 0\right)$, y -intercept $\left(0, \frac{C}{B}\right)$ **1.5.6.21.****Answer.**

a $a = 100 + 150t$

- b The slope tells us that the skier's altitude is increasing at a rate of 150 feet per minute, the vertical intercept that the skier began at an altitude of 200 feet.

1.5.6.23.**Answer.**

a $G = 25 + 12.5t$

- b The slope tells us that the garbage is increasing at a rate of 12.5 tons per year, the vertical intercept that the dump already had 25 tons (when the new regulations went into effect).

1.5.6.25.**Answer.**

a $M = 7000 - 400w$


- b The slope tells us that Tammy's bank account is diminishing at a rate of \$400 per week, the vertical intercept that she had \$7000 (when she lost all sources of income).

1.5.6.27.

Answer.

a 50°F

b -20°C




d The slope, $\frac{9}{5} = 1.8$, tells us that Fahrenheit temperatures increase by 1.8° for each increase of 1° Celsius.

e C -intercept $(-17\frac{7}{9}, 0)$: $-17\frac{7}{9}^{\circ}\text{C}$ is the same as 0°F ; F -intercept $(0, 32)$: 0°C is the same as 32°F .

1.5.6.29.

Answer.




b $m = 25, b = 250$

c $y = 250 + 25x$


1.5.6.31.

Answer.




b $y = 0.12x + 25.4$

c 18 kg

1.5.6.33.**Answer.**

b $y + 5 = -3(x - 2)$

c $y = 1 - 3x$

1.5.6.35.**Answer.**


b $y + 1 = \frac{5}{3}(x - 2)$

c $y = \frac{-13}{3} + \frac{5}{3}x$

1.5.6.37.**Answer.**


a $y + 3.5 = -0.25(x + 6.4)$

b $y = -5.1 - 0.25x$

**1.5.6.39.****Answer.**

a $y + 250 = 2.4(x - 80)$

b $y = -442 + 2.4x$

**1.5.6.41.****Answer.**

a $m = \frac{2}{3}$

b $y = \frac{-1}{3} + \frac{2}{3}x$

1.5.6.43.**Answer.**

a $(-4, 4)$: neither; $(0, 3)$:
 $y = px + q$; $(3, 2)$: both; $(2, 1)$:
 neither; $(1, -2)$: $y = tx + v$

b $p = \frac{-1}{3}$, $q = 3$, $t = 2$, $v = -4$

1.5.6.45.**Answer.**

a $m = 4$, $b = 40$

b $y = 40 + 4x$

1.5.6.47.**Answer.**

a $m = -80$, $b = -2000$

b $P = -2000 - 80t$


1.5.6.49.**Answer.**

a $m = \frac{1}{4}, b = 0$

b $V = \frac{1}{4}d$

1.5.6.51.**Answer.**

a $y = \frac{3}{4}x, y = 1 + \frac{3}{4}x, y = -2.7 + \frac{3}{4}x$



The lines are parallel.

1.5.6.53.**Answer.**

a II

b III

c I

d IV


1.5.6.55.**Answer.**

a III

b IV

c II

d I

1.5.6.57.**Answer.** $m = 2; (6, -1)$ **1.5.6.59.****Answer.** $m = \frac{-4}{3}; (-5, 3)$ **1.5.6.61.****Answer.**b The lines with slope 3 and $\frac{-1}{3}$ are perpendicular to each other, and the lines with slope -3 and $\frac{1}{3}$ are perpendicular to each other.**1.5.6.63.****Answer.** $m = -0.0018$ degree/foot, so the boiling point drops with altitude at a rate of 0.0018 degree per foot. $b = 212$, so the boiling point is 212° at sea level (where the elevation $h = 0$).

1.6 · Chapter Summary and Review

1.6.2 · Chapter 1 Review Problems

1.6.2.1.


Answer.

a

n	100	500	800	1200	1500
C	4000	12,000	18,000	26,000	32,000

b $C = 20n + 2000$

c



d \$22,000

e 400

1.6.2.3.

Answer.

a $R = 2100 - 28t$

b $(75, 0)$, $(0, 2100)$


c t -intercept: The oil reserves will be gone in 2080; R -intercept: There were 2100 billion barrels of oil reserves in 2005.

1.6.2.5.



Answer.

a $2C + 5A = 1000$

b $(500, 0)$, $(0, 200)$



c C -intercept: If no adult tickets are sold, he must sell 500 children's tickets; A -intercept: If no children's tickets are sold, he must sell 200 adult tickets.

1.6.2.7.**Answer.****1.6.2.11.****Answer.****1.6.2.15.****Answer.** A function: Each x has exactly one associated y -value.**1.6.2.19.****Answer.** $N(10) = 7000$: Ten days after the new well is opened, the company has pumped a total of 7000 barrels of oil.**1.6.2.21.****Answer.**
Function**1.6.2.23.****Answer.** Not
a function**1.6.2.25.****Answer.** $F(0) = 1$, $F(-3) = \sqrt{37}$ **1.6.2.27.****Answer.** $h(8) = -6$, $h(-8) = -14$ **1.6.2.29.****Answer.**

a $f(-2) = 3$, $f(2) = 5$

b $t = 1$, $t = 3$

c t -intercepts $(-3, 0)$, $(4, 0)$; $f(t)$ -intercept: $(0, 2)$

d Maximum value of 5 occurs at $t = 2$



1.6.2.31.**Answer.**

a $x = \frac{1}{2} = 0.5$

c $x > 4.9$

b $x = \frac{27}{8} \approx 3.4$

d $x \leq 2.0$

1.6.2.33.**Answer.****1.6.2.9.****Answer.****1.6.2.13.****Answer.**

a $x \approx \pm 5.8$

c $-2.5 < x < 0$ or $0 < x < 2.5$

b $x = \pm 0.4$

d $x \leq -0.5$ or $x \geq 0.5$

1.6.2.35.

Answer. $H(2a) = 4a^2 + 4a$, $H(a+1) = a^2 + 4a + 3$


1.6.2.37.

Answer. $f(a) + f(b) = 2a^2 + 2b^2 - 8$, $f(a+b) = 2a^2 + 4ab + 2b^2 - 4$

1.6.2.39.**Answer.** The volleyball**1.6.2.41.****Answer.** Highway 33**1.6.2.43.****Answer.**


a $B = 800 - 5t$

b

c $m = -5$ thousand barrels/minute: The amount of oil in the tanker is decreasing by 5000 barrels per minute.**1.6.2.45.****Answer.**

a $F = 500 + 0.10C$

b

c $m = 0.10$: The fee increases by \$0.10 for each dollar increase in the remodeling job.**1.6.2.47.**

Answer. $\frac{-3}{2}$

1.6.2.49.

Answer. $\frac{-34}{83} \approx -0.4$

1.6.2.51.**Answer.** 80 ft**1.6.2.53.****Answer.**

a $h(x_2) - h(x_1)$

b $\frac{h(x_2) - h(x_1)}{x_2 - x_1}$

1.6.2.55.

Answer. Neither


1.6.2.57.

Answer.

d	V
-5	-4.8
-2	-3
	-1.2
6	1.8
10	4.2

1.6.2.59.Answer. $m = \frac{1}{2}, b = \frac{-5}{4}$ **1.6.2.61.**Answer. $m = -4, b = 3$ **1.6.2.63.**

Answer.



b $y = \frac{10}{3} - \frac{2}{3}x$

1.6.2.65.

Answer.

a $m = -2, b = 3$

b $y = 3 - 2x$

1.6.2.67.Answer. $\frac{3}{5}$ **1.6.2.69.**

Answer.

a $\frac{3}{2}$

b (4, 2), no

c (6, 5)

1.6.2.71.

Answer. (3, -14), (-7, 2)

1.6.2.73.

Answer.

a $T = 62 - 0.0036h$

b $-46^\circ\text{F}; 108^\circ\text{F}$

c -71°F

1.6.2.75.Answer. $y = \frac{2}{5} - \frac{9}{5}x$

1.6.2.77.**Answer.**


a

t	0	15
P	4800	6780

b $P = 4800 + 132t$

c $m = 132$ people/year: the population grew at a rate of 132 people per year.**1.6.2.79.****Answer.** 6**1.6.2.81.****Answer.**

a



b 129 lb, 145 lb

c $y = 2.6x - 44.3$


d 137 lb

e $y = 2.84x - 55.74$

f 137.33 lb

1.6.2.83.**Answer.**

a



b 45 cm

c 87 cm

d $y = 1.2x - 3$

e 69 cm

f $y = 1.197x - 3.660$; 68.16 cm**2 · Modeling with Functions****2.1 · Nonlinear Models****2.1.7 · Nonlinear Models (Homework 2.1)****2.1.7.1.**Answer. $\pm\frac{5}{3}$ **2.1.7.3.**Answer. $\pm\sqrt{6}$ **2.1.7.5.**Answer. $\pm\sqrt{6}$ **2.1.7.7.**Answer. ± 2.65 **2.1.7.9.**Answer. ± 5.72 **2.1.7.11.**Answer. ± 5.73 **2.1.7.13.**Answer. $\pm\sqrt{\frac{Fr}{m}}$ **2.1.7.15.**Answer. $\pm\sqrt{\frac{2s}{g}}$ **2.1.7.17.**


Answer.

a $V = 2.8\pi r^2 \approx 8.8r^2$

r	1	2	3	4	5	6	7	8
V	8.8	35.2	79.2	140.7	219.9	316.7	431.0	563.0

b The volume increases by a factor of 4.

c 5.86 cm


**2.1.7.19.**

Answer. 21 in.

2.1.7.21.Answer. $\sqrt{1800} \approx 42.4$ m**2.1.7.23.**Answer. $\sqrt{128}$ in. by $\sqrt{128}$ in. ≈ 11.3 in. $\times 11.3$ in.

2.1.7.25.


Answer.



a
b $x = \pm 12$

2.1.7.27.


Answer.



a
b $x = 1$ or $x = 9$

2.1.7.29.

Answer.



a
b $x = 10$ or $x = -2$

2.1.7.31.

Answer. 5, -1

2.1.7.33.

Answer. $\frac{5}{2}, \frac{-3}{2}$

2.1.7.35.

Answer. $-2 \pm \sqrt{3}$

2.1.7.37.

Answer. $\frac{1}{2} \pm \frac{\sqrt{3}}{2}$

2.1.7.39.

Answer. $\frac{-2}{9}, \frac{-4}{9}$

2.1.7.41.

Answer. $\frac{7}{8} \pm \frac{\sqrt{8}}{8}$

2.1.7.43.

Answer. 6

2.1.7.45.

Answer. 64

2.1.7.47.

Answer. $\frac{13}{6}$

2.1.7.49.

Answer. 8

2.1.7.51.

Answer. 9


2.1.7.53.

Answer. $\frac{33}{64}$

2.1.7.55.

Answer.

a $B = 5000(1+r)^2$



b 11.8%

2.1.7.57.

Answer. 8%


2.1.7.59.

Answer. 7.98 mm

2.1.7.61.

Answer.

a $\sqrt{3} \approx 1.73$ sq cm, $4\sqrt{3} \approx 6.93$ sq cm, $25\sqrt{3} \approx 43.3$ sq cm



c An equilateral triangle with side 5.1 cm has area 11.263 cm².

d side ≈ 6.8 cm

e $\frac{\sqrt{3}}{4}s^2 = 20$; $s \approx 6.8$

f ≈ 20 cm

2.1.7.63.

Answer. $\pm\sqrt{\frac{bc}{a}}$

2.1.7.65.

Answer. $a \pm 4$


2.1.7.67.

Answer. $\frac{-b \pm 3}{a}$

2.1.7.69.

Answer.

Height	Base	Area	Height	Base	Area
1	34	34	10	16	160
2	32	64	11	14	154
3	30	90	12	12	144
4	28	112	13	10	130
5	26	130	14	8	112
6	24	144	15	6	90
7	22	154	16	4	64
8	20	160	17	2	34
9	18	162	18	0	0



c 162 sq ft, with base 18 ft, height 9 ft

d Base: $36 - 2x$; area: $x(36 - 2x)$

e See (a)


f 6.5 ft or 11.5 ft

2.1.7.71.

Answer.

a

v	0	1	2	3	4	5	6	7	8	9	10	11
J	0	0.05	0.2	0.46	0.82	1.28	1.84	2.5	3.27	4.13	5.1	6.17



c	v	0	1	2	3	4	5	6	7	8	9	10	11
	H	0.9	0.95	1.1	1.36	1.72	2.18	2.74	3.4	4.17	5.03	6.0	7.07

d 5.5 meters

e 10.15 meters per second

2.2 · Some Basic Functions

2.2.7 · Some Basic Functions (Homework 2.2)

2.2.7.1.

Answer.

a -9

b 9

2.2.7.5.

Answer. -50

2.2.7.9.

Answer. 1

2.2.7.11.

Answer.

a 2.7

b -2.7

c 1.8

d 2.9

2.2.7.13.

Answer.


a 0.3

b -0.4

c 0.2

2.2.7.15.

Answer.



b $x = 0, x = 1$

c $(-\infty, 0)$ and $(0, 1)$

2.2.7.3.

Answer.

a -4


b 20

2.2.7.7.

Answer. 144

2.2.7.17.

Answer.



b $x = 1$

c $(1, +\infty)$

2.2.7.19.

Answer. Graph (b) is the basic graph shifted 2 units down; graph (c) is the basic graph shifted 1 unit up.

2.2.7.23.

Answer. Graph (b) is the basic graph reflected about the x -axis; graph (c) is the basic graph reflected about the y -axis.

2.2.7.21.

Answer. Graph (b) is the basic graph shifted 1.5 units left; graph (c) is the basic graph shifted 1 unit right.

2.2.7.25.**Answer.**

$$\begin{array}{lll} \text{a } \sqrt{x} & \text{c } |x| & \text{e } x^3 \\ \text{b } \sqrt[3]{x} & \text{d } \frac{1}{x} & \text{f } \frac{1}{x^2} \end{array}$$

2.2.7.27.**Answer.**

$$\text{a } x \approx 12 \qquad \text{b } x \approx 18 \qquad \text{c } x < 9 \qquad \text{d } x > 3$$

2.2.7.29.**Answer.**

$$\text{a } t \approx -3.1 \qquad \text{b } t \approx 1.5 \qquad \text{c } t < 0.8 \qquad \text{d } -2.4 < t < 0.4$$


2.2.7.31.**Answer.**

$$\text{a } x = 41 \qquad \text{b } 29 < x < 61$$

2.2.7.33.**Answer.**


$$\text{a } x = -5 \text{ or } x = 17 \qquad \text{b } -1 < x < 13$$

2.2.7.35.**Answer.**

$$\text{a } \begin{array}{|c|c|c|c|c|c|c|c|} \hline x & 4 & 1 & \frac{1}{4} & 0 & \frac{1}{4} & 1 & 4 \\ \hline y & -2 & -1 & -\frac{1}{2} & 0 & \frac{1}{2} & 1 & 2 \\ \hline \end{array}$$



b no

2.2.7.37.**Answer.**

$$\text{a } \begin{array}{|c|c|c|c|c|c|c|c|} \hline x & 2 & 1 & \frac{1}{2} & 0 & \frac{1}{2} & 1 & 2 \\ \hline y & -2 & -1 & -\frac{1}{2} & 0 & \frac{1}{2} & 1 & 2 \\ \hline \end{array}$$


b no


2.2.7.39.**Answer.**



b yes


2.2.7.41.

Answer.




2.2.7.45.

Answer.



2.2.7.49.


Answer.



2.2.7.53.


Answer.

$$f(x) = \begin{cases} 8 - 2x & x < 4 \\ 2x - 8 & x \geq 4 \end{cases}$$




2.2.7.43.

Answer.




2.2.7.47.

Answer.



2.2.7.51.


Answer.



2.2.7.55.


Answer.

$$g(t) = \begin{cases} -1 - \frac{t}{3} & t < -3 \\ 1 + \frac{t}{3} & t \geq -3 \end{cases}$$



2.2.7.57.

$$\text{Answer. } F(x) = \begin{cases} -x^3 & x < 0 \\ x^3 & x \geq 0 \end{cases}$$



2.2.7.59.

Answer.

a Not always true:

$$f(1+2) \neq f(1) + f(2)$$

because $9 \neq 5$.

b True: $(ab)^2 = a^2b^2$

2.2.7.63.

Answer.

a Not always true (unless

$$b = 0): f(1+2) \neq f(1) + f(2)$$

because $3m + b \neq 3m + 2b$.





b Not always true:

$$f(1 \cdot 2) \neq f(1) \cdot f(2) \text{ because}$$

$$2m + b \neq 2m^2 + 3mb + b^2.$$

2.2.7.65.

Answer.



2.2.7.67.

Answer. The distributive law shows a relationship between multiplication and addition that always holds. The equation $f(a+b) = f(a) + f(b)$ is not about multiplication and may or may not be true.

2.3 · Transformations of Graphs

2.3.5 · Transformations of Graphs (Homework 2.3)

2.3.5.1.

$$\text{Answer. } y = \sqrt{x+2}$$

2.3.5.5.


$$\text{Answer. } y = \frac{1}{x-4}$$

2.3.5.3.


$$\text{Answer. } y = x^3 - 1$$

2.3.5.7.**Answer.**

- a Translate $y = |x|$ by 2 units down.

**2.3.5.13.****Answer.**


- a Translate $y = r^3$ by 2 units left.

**2.3.5.19.**


Answer. A vertical stretch by a factor of 3: $y = \frac{3}{x}$

2.3.5.23.**Answer.**


- a Scale factor $\frac{1}{3}$; $y = |x|$ is compressed vertically by the scale factor.

**2.3.5.29.****Answer.**


- a Scale factor $\frac{-1}{2}$; $y = s^3$ is reflected over the s -axis and compressed vertically by a factor of $\frac{1}{2}$.

**2.3.5.33.****Answer.****2.3.5.9.****Answer.**

- a Translate $y = \sqrt[3]{s}$ by 4 units right.

**2.3.5.15.****Answer.**


- a Translate $y = \sqrt{d}$ by 3 units down.

**2.3.5.21.**


Answer. A vertical compression, the scale factor is $\frac{1}{2}$: $y = \frac{1}{2}x^3$

2.3.5.25.**Answer.**


- a Scale factor -2 ; $y = \frac{1}{z^2}$ is reflected over the z -axis and stretched vertically by a factor of 2.

**2.3.5.31.****Answer.**


- a Scale factor $\frac{1}{3}$; $y = \frac{1}{x}$ is compressed vertically by the scale factor.

**2.3.5.11.****Answer.**


- a Translate $y = \frac{1}{t^2}$ by 1 unit up.

**2.3.5.17.****Answer.**

- a Translate $y = \frac{1}{v}$ by 6 units left.

**2.3.5.27.****Answer.**

- a Scale factor -3 ; $y = \sqrt{v}$ is reflected over the v -axis and stretched vertically by a factor of 3.



- | | | |
|------|------|-------|
| a vi | c iv | e v |
| b ii | d i | f iii |

2.3.5.35.**Answer.**

- a Vertical stretch by a factor of 3: $y = 3f(x)$
- b Reflection about the x -axis: $y = -f(x)$
- c Translation 1 unit right: $y = f(x - 1)$
- d Translation 4 units up: $y = f(x) + 4$

2.3.5.37.**Answer.**

- a Reflection about the v -axis and vertical stretch by a factor of 2:
 $T = -2h(v)$
- b Vertical stretch by a factor of 3: $T = 3h(v)$
- c Translation 3 units up: $T = h(v) + 3$
- d Translation 3 units left: $T = h(v + 3)$

2.3.5.39.**Answer.**

- a Translation 2 units up: $y = f(x) + 2$
- b Translation 4 units down: $y = f(x) - 4$
- c Vertical compression by a factor of $\frac{1}{2}$: $y = \frac{1}{2}f(x)$
- d Translation 1 unit right: $y = f(x - 1)$

2.3.5.41.**Answer.**

- a Translation 1 unit right: $y = f(x - 1)$
- b Part (a) is translated 30 units up: $y = f(x - 1) + 30$
- c f is reflected about the x -axis and stretched vertically by a factor of 2: $y = -2f(x)$
- d Part (c) is translated 10 units down: $y = -2f(x) - 10$

2.3.5.43.**Answer.**

$y = \frac{1}{2} \cdot \frac{1}{x^2}$ is a vertical compression with factor $\frac{1}{2}$ of $y = \frac{1}{x^2}$.

2.3.5.45.**Answer.**

$y = 2\sqrt[3]{x}$ is a vertical stretch with factor 2 of $y = \sqrt[3]{x}$.

2.3.5.47.**Answer.**

$y = 3|x|$ is a vertical stretch with factor 3 of $y = |x|$.


2.3.5.49.**Answer.**

$y = \frac{1}{8}x^3$ is a vertical compression with factor $\frac{1}{8}$ of $y = x^3$.

2.3.5.51.**Answer.**


- a Translation by 2 units up and 3 units right

b

**2.3.5.55.****Answer.**


- a Reflection across the u -axis, vertical stretch by a factor of 3, translation by 4 units left and 4 units up

b

**2.3.5.59.****Answer.**

- a Reflection across the w -axis, vertical stretch by a factor of 2, translation by 6 units up and 1 unit right

b

**2.3.5.63.****Answer.**

- a Translation by 4 units up and 1 unit right: $y = f(x - 1) + 4$
- b Vertical stretch by a factor of 2 and a translation by 4 units up: $y = 2f(x) + 4$


2.3.5.65.**Answer.**

- a $y = |x|$ translated by 1 unit left and 2 units down
- b $y = |x + 1| - 2$

2.3.5.53.**Answer.**


- a Translation by 2 units left and 3 units down.

b

**2.3.5.57.****Answer.**


- a Vertical stretch by a factor of 2, translation by 5 units right and 1 down

b

**2.3.5.61.****Answer.**

- a Translation by 8 units right and 1 unit down

b

**2.3.5.67.****Answer.**

- a $y = \sqrt{x}$ reflected about the x -axis and shifted 3 units up
- b $y = -\sqrt{x} + 3$

2.3.5.69.**Answer.**

a $y = x^3$ translated by 3 units
right and 1 unit up

b $y = (x - 3)^3 + 1$

2.3.5.71.**Answer.**

a $y = f(x - 20)$: Students scored 20 points higher than Professor Hilbert's class.

b $y = 1.5f(x)$: The class is about 50% larger than Hilbert's, but the classes scored the same.

2.3.5.73.**Answer.**



a $y = f(x - 5000)$: Taxpayers earn \$5000 more than Californians in each tax rate

b $y = f(x) - 0.2$: Taxpayers pay 0.2% less tax than Californians on the same income.

2.3.5.75.**Answer.**

a $y = g(t + 2)$: This population has its maximum and minimum two months before the marmots.

b $y = g(t) - 20$: This population remains 20 fewer than that of the marmots.

2.4 · Functions as Mathematical Models**2.4.5 · Functions as Mathematical Models (Homework 2.4)****2.4.5.1.****Answer.** (b)**2.4.5.3.****Answer.** (a)**2.4.5.5.****Answer.****2.4.5.7.****Answer.**

2.4.5.9.

Answer. (b)

2.4.5.11.

Answer.

a II


b IV

c I

d III


2.4.5.13.

Answer.




2.4.5.15.

Answer.




2.4.5.17.


Answer.



2.4.5.19.


Answer. $y = x^3$ stretched or compressed vertically

2.4.5.21.

Answer. $y = \frac{1}{x}$ stretched or compressed vertically

2.4.5.23.

Answer. $y = \sqrt{x}$

**2.4.5.25.****Answer.**

- a Increasing
- b Concave up

2.4.5.27.**Answer.**

- a Increasing
- b Concave down

2.4.5.29.**Answer.**


- a Increasing,
linear (neither
concave up nor
down)
- b C

2.4.5.31.**Answer.**


- a Increasing,
concave down
- b F

2.4.5.33.**Answer.**


- a Decreasing,
linear (neither
concave up nor
down)
- b D

2.4.5.35.**Answer.**

$$y = 4\sqrt[3]{x}$$

2.4.5.37.**Answer.**

$$y = 3 \cdot \frac{1}{x^2}$$

2.4.5.39.**Answer.**

$$y = 0.5x^2$$

2.4.5.41.**Answer.**

- a Table (4), Graph (C)
- b Table (3), Graph (B)

- c Table (1), Graph (D)
- d Table (2), Graph (A)


2.4.5.43.**Answer.**

- a III

- b 3


2.4.5.45.**Answer.**

$$\text{a } S(x) = \begin{cases} 5.95 & x \leq 25 \\ 7.95 & 25 < x \leq 50 \\ 9.95 & 50 < x \leq 75 \\ 10.95 & 75 < x \leq 100 \end{cases}$$



2.4.5.47.

Answer.




During the first 400 seconds Bob's altitude is climbing with the aircraft; then the aircraft maintains a constant altitude of 10,000 feet for the next 100 seconds; after jumping from the plane, Bob falls for 20 seconds before opening the parachute; he falls at a constant rate after the chute opens.

b 240 seconds (4 minutes) and $500 + \sqrt{250} \approx 515.8$

2.4.5.49.

Answer.



b $m \approx 3.2$ mm/cc: The height of precipitate increases by 1 mm for each additional cc of lead nitrate

$$\text{c } f(x) = \begin{cases} 1.34 + 3.2x & x < 2.6 \\ 9.6 & x \geq 2.6 \end{cases}$$

d The increasing portion of the graph corresponds to the period when the reaction was occurring, and the horizontal section corresponds to when the potassium iodide is used up.

2.4.5.51.

Answer.

a II

b IV

c I

d III

2.5 · The Absolute Value Function


2.5.7 · The Absolute Value Function (Homework 2.5)

2.5.7.1.

Answer.

a $|x| = 6$

b




2.5.7.3.

Answer.

a $|p + 3| = 5$

b




2.5.7.5.

Answer.

a $|t - 6| < 3$

b




2.5.7.7.

Answer.


a $|b + 1| \geq 0.5$

b



2.5.7.9.

Answer.




a $x = -5$ or $x = -1$

b $-7 \leq x \leq 1$

c $x < -8$ or $x > 2$

2.5.7.11.

Answer.



a $x = 4$

b No solution

c No solution

2.5.7.13.

Answer. $x = \frac{-3}{2}$ or
 $x = \frac{5}{2}$

2.5.7.15.

Answer. $q = \frac{-7}{3}$

2.5.7.17.

Answer. $b = -14$ or
 $b = 10$

2.5.7.19.

Answer. $w = \frac{13}{2}$ or
 $w = \frac{15}{2}$

2.5.7.21.

Answer. No
solution

2.5.7.23.

Answer. No
solution

2.5.7.25.

Answer.
 $\frac{-9}{2} < x < \frac{-3}{2}$

2.5.7.27.

Answer. $d \leq -2$ or
 $d \geq 5$

2.5.7.29.

Answer. All real
numbers

2.5.7.31.

Answer.
 $1.4 < t < 1.6$

2.5.7.33.

Answer. $T \leq 3.2$
or $T \geq 3.3$

2.5.7.35.

Answer. No
solution

2.5.7.37.

Answer. $4.299 < l < 4.301$

2.5.7.39.

Answer. $250 \leq t \leq 350$

2.5.7.41.


Answer. $|T - 5| < 0.3$

2.5.7.43.

Answer. $|D - 100| \leq 5$

2.5.7.45.

Answer. $|g - 0.25| \leq 0.001$



2.5.7.47.

Answer.

a $|t - 200| < 50$, $150 \leq t < 250$

b $|t - 200| < 0.5$, $199.5 \leq t < 200.5$

c $|t - 200| < 0.05$, $199.95 \leq t < 200.05$

2.5.7.49.**Answer.**

$$\text{a } |3x - 6| = \begin{cases} -(3x - 6) & \text{if } x < 2 \\ 3x - 6 & \text{if } x \geq 2 \end{cases}$$

$$\text{b } -(3x - 6) \leq 9, \quad 3x - 6 < 9$$

$$\text{c } -1 < x < 5$$

d The solutions are the same.


2.5.7.51.**Answer.**

$$\text{a } |2x + 5| = \begin{cases} -(2x + 5) & \text{if } x < -\frac{5}{2} \\ 2x + 5 & \text{if } x \geq -\frac{5}{2} \end{cases}$$

$$\text{b } -(2x + 5) > 7, \quad 2x + 5 > 7$$

$$\text{c } x < -6 \text{ or } x > 1$$

d The solutions are the same.


2.5.7.53.**Answer.**

$$\text{a } f(x) = \begin{cases} -2x, & x < -4 \\ 8, & -4 \leq x \leq 4 \\ 2x, & x > 4 \end{cases}$$

b The graph looks like a trough. The middle horizontal section is $y = p + q$ for $-p \leq x \leq q$, the left side, $x < -p$, has slope -2 and the right side, $x > q$, has slope 2 .

$$\text{c } g(x) = \begin{cases} -2x + q - p, & x < -p \\ p + q, & -p \leq x \leq q \\ 2x + p - q, & x > q \end{cases}$$

2.5.7.55.**Answer.**



$$\text{a } f(x) = \begin{cases} -3x, & x < -4 \\ -x + 8, & -4 \leq x \leq 0 \\ x + 8, & 0 < x < 4 \\ 3x, & x \geq 4 \end{cases}$$

b 8

c $p + q$


2.5.7.57.

Answer.

a $|x + 12|, |x + 4|, |x - 24|$

b $f(x) = |x + 12| + |x + 4| + |x - 24|$

c



At x -coordinate -4

2.5.7.59.

Answer. 2 miles east of the river

2.6 · Domain and Range

2.6.6 · Domain and Range (Homework 2.6)

2.6.6.1.

Answer. Domain: $[-5, 3]$; Range: $[-3, 7]$

2.6.6.5.

Answer. Domain: $[-2, 2]$; Range: $[-1, 1]$

2.6.6.3.

Answer. Domain: $[-4, 5]$; Range: $[-1, 1) \cup [3, 6]$

2.6.6.7.

Answer. Domain: $(-5, 5]$; Range: $\{-1, 0, 2, 3\}$

2.6.6.9.**Answer.**

- a Domain: all real numbers;
Range: all real numbers
- b Domain: all real numbers;
Range: $[0, \infty)$

2.6.6.11.**Answer.**


- a Domain: all real numbers
except zero; Range: $(0, \infty)$
- b Domain: all real numbers
except zero; Range: all real
numbers except zero

2.6.6.13.**Answer.** Domain: $[0, 26.2]$; Range: $[90, 300]$ **2.6.6.15.****Answer.** Domain: $[0, 600]$; Range: $[-90, 700]$ **2.6.6.17.****Answer.**

- a $V(t) = 6000 - 550t$
- b Domain: $[0, 10]$; Range: $[500, 6000]$

2.6.6.19.**Answer.**


a



- b Domain: $[0, 4]$; Range: $[0, 64]$. The ball reaches a height of 64 feet and hits the ground 4 seconds after being hit.

2.6.6.21.**Answer.**

a





- b Range: $\{2.50, 2.90, 3.30, 3.70, 4.10\}$
- c \$13.30

2.6.6.23.

Answer. Domain: nonnegative integers; The range includes all whole number multiples of 2.50 up to $20 \times 2.50 = 50$, all integer multiples of 2.25 from $21 \times 2.25 = 47.25$ to $50 \times 2.25 = 112.50$ and all integer multiples of 2.10 from $51 \times 2.10 = 107.10$ onwards: 0, 2.50, 5.00, 7.50, ..., 50, 47.25, 49.50, 51.75, ..., 112.50, 107.10, 109.20, 111.30, ...



2.6.6.25.

Answer.

a $f(x)$ domain: $x \neq 4$; Range: $(0, \infty)$ b $h(x)$ domain: $x \neq 0$; Range: $(-4, \infty)$ 


2.6.6.27.


Answer.

a $G(v)$ domain: all real numbers; Range: all real numbersb $H(v)$ domain: all real numbers; Range: all real numbers

2.6.6.29.

Answer.

a $G(v)$ domain: $[2, \infty)$; Range: $[0, \infty)$ b $H(v)$ domain: $[0, \infty)$; Range: $[-2, \infty)$

**2.6.6.31.****Answer.**

a Not in range

b $x = -6$ or $x = 2$ **2.6.6.35.****Answer.**a $w = \frac{1}{2}$

b Not in range

2.6.6.33.**Answer.**a $t = -64$ b $t = -8$ **2.6.6.37.****Answer.**


a Not in range

b $h = 4$ **2.6.6.39.****Answer.** Domain: $[-2, 5]$; Range: $[-4, 12]$ **2.6.6.43.****Answer.** Domain: $[-2, 2]$; Range: $[-9, 7]$ **2.6.6.47.****Answer.** Domain: $[-1.25, 2.75]$; Range: $\left[\frac{4}{17}, 4\right]$ **2.6.6.41.****Answer.** Domain: $[-5, 3]$; Range: $[-15, 1]$ **2.6.6.45.****Answer.** Domain: $[-1, 8]$; Range: $[0, 3]$ **2.6.6.49.****Answer.** Domain: $(3, 6]$; Range: $\left[-\infty, \frac{-1}{3}\right]$ **2.6.6.51.****Answer.**

a Squaring both sides of the equation gives the equation of the circle centered on the origin with radius 4, but the points in the third and fourth quadrants are extraneous solutions introduced by squaring. (The original equation allowed only $y \geq 0$.)

b Domain: $[-4, 4]$; Range: $[0, 4]$

c



The calculator does not show the graph extending down to the x -axis.

2.6.6.53.**Answer.**a Domain: $x \neq 2$; Range: $(0, \infty)$ b Domain: $x \neq 0$; Range: $(-2, \infty)$ c Domain: $x \neq 3$; Range: $(-5, \infty)$ **2.6.6.57.****Answer.**a Domain: $[3, 13]$; Range: $[-2, 2]$ b Domain: $[0, 10]$; Range: $[-6, 6]$ c Domain: $[5, 15]$; Range: $[-4, 4]$ **2.6.6.61.****Answer.**a $f(x)$ b $g(x)$ **2.6.6.63.****Answer.**a $g(x)$ b $f(x)$ **2.6.6.65.****Answer.** Domain: $[0^\circ, 90^\circ]$; Range: $[12, 24]$

2.7 · Chapter Summary and Review

2.7.2 · Chapter 2 Review Problems

2.7.2.1.**Answer.** $x = 1$ or $x = 4$ **2.7.2.3.****Answer.** $w = -2$ or $w = 4$ **2.7.2.5.****Answer.** $r = -1 \pm \sqrt{\frac{A}{P}}$ **2.7.2.7.****Answer.** 11%**2.7.2.9.****Answer.** $P = 1.001$ **2.7.2.11.****Answer.** $m = 29$ **2.7.2.13.****Answer.** $r = 3$ **2.7.2.15.****Answer.** $\sqrt{12,132} \approx 110$ cm

2.7.2.17.Answer. -2 **2.7.2.19.**Answer. 24 **2.7.2.21.**

Answer.

a $x = -2$ or $x = 6$

b $(-2, 6)$

c $(-\infty, -2] \cup [6, +\infty)$

2.7.2.23.

Answer.


a $x = 0$ or $x = 3$

b $(-\infty, 0) \cup (3, \infty)$


c $(0, 3)$

2.7.2.25.


Answer.

**2.7.2.27.**

Answer.


**2.7.2.29.**

Answer.

**2.7.2.31.**


Answer.

a $y = |x|$ shifted up 2 units

**2.7.2.33.**


Answer.

a $y = \sqrt{x}$ shifted up 3 units




2.7.2.35.**Answer.**

- a $y = |x|$ shifted left 2 units and down 3 units

**2.7.2.37.****Answer.**

- a $y = \sqrt{x}$ reflected across the horizontal axis and stretched vertically by a factor of 2

**2.7.2.39.****Answer.**

a $y = \frac{-3}{2}f(t)$

b $y = \frac{-3}{2}f(t) + 3$


c $y = \frac{-3}{2}f(t+2) + 3$

2.7.2.41.**Answer.**


a $y = f(t-1)$

b $y = -f(t-1)$

c $y = -f(t-1) + 300$

2.7.2.43.**Answer.** $y = (x-2)^2 - 4$ **2.7.2.45.****Answer.****2.7.2.47.****Answer.** I (c), II (b), III (a)**2.7.2.49.**

$$\text{Answer. } g(t) = \begin{cases} 60 + t, & 0 \leq t < 10 \\ 70, & 10 \leq t < 30 \\ 70 - \frac{1}{2}(t - 30), & 30 \leq t \leq 60 \end{cases}$$



2.7.2.51.

$$\text{Answer. } |x| = 4$$

2.7.2.53.

$$\text{Answer. } |p - 7| < 4$$

2.7.2.55.

$$\text{Answer. } t = \frac{6}{5} \text{ or } t = \frac{12}{5}$$

2.7.2.57.

Answer. No solutions

2.7.2.59.

$$\text{Answer. } p = 1 \text{ or } p = 6$$

2.7.2.61.

$$\text{Answer. } \left(\frac{-2}{3}, 2 \right)$$

2.7.2.63.

$$\text{Answer. } (-\infty, -0.9] \cup [0.1, \infty)$$

2.7.2.65.


$$\text{Answer. } |H - 65.5| < 9.5$$

2.7.2.67.

$$\text{Answer. } [2.05, 2.15]$$

2.7.2.69.

Answer.



b $g(x) = \frac{24}{x}$

2.7.2.71.

Answer.

a

x	0	4	8	14	16	22
y	24	20	16	10	8	2

b $y = 24 - x$

2.7.2.73.

Answer.

a

x	0	1	4	9	16	25
y	0	1	2	3	4	5

b $y = \sqrt{x}$

2.7.2.75.

Answer.

a

x	-3	-2	-1	0	1	2
y	5	0	-3	-4	-3	0

b $y = x^2 - 4$

2.7.2.77.

Answer. Domain: $[-2, 4]$; Range: $[-10, -4]$

2.7.2.79.

Answer. Domain: $(-2, 4]$; Range: $\left[\frac{1}{6}, \infty\right)$

3 · Power Functions

3.1 · Variation

3.1.9 · Variation (Homework 3.1)

3.1.9.1.

Answer.


a

Price of item	18	28	12
Tax	1.17	1.82	0.78
Tax/Price	0.065	0.065	0.065

Yes; 6.5%

b $T = 0.065p$

c



3.1.9.3.

Answer.


a

Width (feet)	2	2.5	3
Length (feet)	12	9.6	8
Length \times width	24	24	24

24 square feet

b $L = \frac{24}{w}$

c

**3.1.9.5.****Answer.**

- a The ratio $\frac{y}{x}$ is a constant.
 b The product xy is a constant.

3.1.9.7.**Answer.**

Length	Width	Perimeter	Area
10	8	36	80
12	8	40	96
15	8	46	120
20	8	56	160

a

b No

c $P = 16 + 2l$

d Yes

e $A = 8l$

3.1.9.9.**Answer.** (b)**3.1.9.11.****Answer.** (c)**3.1.9.13.****Answer.**

a $m = 0.165w$

w	50	100	200	400
m	8.25	16.5	33	66

b 19.8 lb

c 303.03 lb

d It will double.

3.1.9.15.**Answer.**

a $L = 0.8125T^2$

T	1	5	10	20
L	0.8125	20.3	81.25	325

b 234.8125 ft

c 0.96 sec

d It must be four times as long.

3.1.9.17.**Answer.**

a $B = \frac{88}{d}$

d	1	2	12	24
B	88	44	7.3	3.7

b 8.8 milligauss

c More than 20.47 in

d It is one half as strong.

3.1.9.19.**Answer.**

a $P = \frac{1825}{8192}w^3 \approx 0.2228w^3$

w	10	20	40	80
P	223	1782	14,259	114,074

b 752 kilowatts

c 33.54 mph

d It is multiplied by 8.

3.1.9.21.**Answer.**

a $y = 0.3x$

x	y
2	0.6
5	1.5
8	2.4
12	3.6
15	4.5

b

c y doubles.**3.1.9.23.****Answer.**

a $y = \frac{2}{3}x^2$

x	y
3	6
6	24
9	
12	96
15	150

b

c y is quadrupled.

3.1.9.25.**Answer.**

a $y = \frac{120}{x}$

	4	30
	8	15
b	20	6
	30	4
	40	3

c y is halved.**3.1.9.27.****Answer.** (b)
 $y = 0.5x^2$ **3.1.9.29.****Answer.** (c)
 $\frac{y}{x^p}$ is not constant for any exponent p .**3.1.9.31.****Answer.** (b)
 $y = \frac{72}{x^2}$ **3.1.9.33.****Answer.** (c)
 $x^p y$ is not constant for any exponent p .**3.1.9.35.****Answer.**

a $d = 0.005v^2$

b 50 m

3.1.9.37.**Answer.**

a $m = \frac{8}{p}$

b 0.8 ton

3.1.9.39.**Answer.**

a $T = \frac{6}{d}$

b 1°C

3.1.9.41.**Answer.**

a $W = 600d^2$

b 864 newtons

3.1.9.43.**Answer.**

a Wind resistance quadruples.

b It is one-ninth as great.

c It is decreased by 19% because it is 81% of the original.

3.1.9.45.**Answer.**

a It is one-fourth the original illumination.

b It is one-ninth the illumination.

c It is 64% of the illumination.

3.1.9.49.

Answer.

a $L = (1.25 \times 10^{-27})m^3$

b 2×10^{12} kg

3.1.9.51.

Answer. $y = kx$ implies that $k(cx) = c(kx) = cy$.

3.1.9.53.

Answer. If $y = kx^2$, then dividing both sides of the equation by x^2 gives $\frac{y}{x^2} = k$.

3.1.9.55.

Answer. Yes

3.2 · Integer Exponents

3.2.6 · Integer Exponents (Homework 3.2)

3.2.6.1.

Answer.

n	-5	-4	-3	-2	-1	0	1	2	3	4	5
3^n	$\frac{1}{243}$	$\frac{1}{81}$	$\frac{1}{27}$	$\frac{1}{9}$	$\frac{1}{3}$	1	3	9	27	81	243

Each time n increases by 1, we multiply the power in the bottom row by 3.

3.2.6.3.

Answer.

a 8 b -8 c $\frac{1}{8}$ d $\frac{-1}{8}$

3.2.6.5.

Answer.

a $\frac{1}{8}$ b $\frac{-1}{8}$ c 8 d -8

3.2.6.7.

Answer.

a $\frac{1}{2^1} = \frac{1}{2}$ b $\frac{1}{(-5)^2} = \frac{1}{25}$ d $\frac{(-2)^4}{16} =$
c $3^3 = 27$

3.2.6.9.

Answer.

a $5 \cdot 4^3 = 320$ $\frac{1}{32q^5}$ c $\frac{-4}{x^2}$ d $8b^3$
b $\frac{1}{(2q)^5} =$

3.2.6.11.**Answer.**

a $\frac{1}{(m-n)^2}$

c $\frac{2p}{q^4}$

b $\frac{1}{y^2} + \frac{1}{y^3}$

d $\frac{-5x^5}{y^2}$

3.2.6.13.**Answer.**

(a)

x	1	2	4	8	16
x^{-2}	1	0.25	0.06	0.02	0.00

(b) The values of $f(x)$ decrease, because x^{-2} is the reciprocal of x^2 .

(c)

x	1	0.5	0.25	0.125	0.0625
x^{-2}	1	4	16	64	256

(d) The values of $f(x)$ increase toward infinity, because x^{-2} is the reciprocal of x^2 .**3.2.6.15.****Answer.** b. (ii), (iii), and (iv) have the same graph, because they represent the same function.**3.2.6.17.****Answer.**

(a) $F(r) = 3r^{-4}$

(b) $G(w) = \frac{2}{5}w^{-3}$

(c) $H(z) = \frac{1}{9}z^{-2}$

3.2.6.19.**Answer.** $x = -1.25$ or $x = 1.25$ **3.2.6.21.****Answer.** $t = \frac{1}{16}$ **3.2.6.23.****Answer.** $v = \frac{1}{5}$ or $v = \frac{-1}{5}$ **3.2.6.25.****Answer.**

(a) $P = 0.355v^3$

(b) $v \approx 52.03$ mph

(c) 3.375

3.2.6.27.**Answer.**

(a) $D = \frac{70}{i}$

(b) It decreases by about 2.3 years.

3.2.6.29.**Answer.**

(a) $L = (4\pi sR^2)T^4 \approx 7.2 \times 10^{-7}R^2T^4$

(b) 4840 K

3.2.6.31.**Answer.**

(a) 500 picowatts


(b) $P = 8000d^{-4}$

(c)


d (nautical miles)	4	5	7	10
P (picowatts)	31.3	12.8	3.3	0.8

(d) 16.8 nautical miles

(e)

**3.2.6.33.****Answer.**(a) $T = 16kr^2$ (b) $T = 0.1r^2$

(c)

**3.2.6.35.****Answer.**(a) a^5 (c) $\frac{1}{p^3}$ (b) $\frac{1}{5^7}$ (d) $\frac{1}{7^{10}}$ **3.2.6.37.****Answer.**(a) $\frac{20}{x^3}$ (b) $\frac{1}{3u^{12}}$ (c) 5^8t **3.2.6.39.****Answer.**(a) $\frac{x^4}{9y^6}$ (b) $\frac{a^6b^4}{36}$ (c) $\frac{5}{6h^6}$

3.2.6.41.

Answer.

(a) $\frac{1}{3}x + 3x^{-1}$

(b) $\frac{1}{4}x^{-2} - \frac{3}{2}x^{-1}$

3.2.6.43.

Answer.

(a) $\frac{1}{2}x^{-2} + x^{-3} - \frac{1}{2}x^{-4}$

(b) $\frac{2}{3}x^{-2} - \frac{1}{9} + \frac{1}{6}x^2$

3.2.6.45.Answer. $x - 3 + 2x^{-1}$ **3.2.6.47.**Answer. $-3 + 6t^{-2} + 12t^{-4}$ **3.2.6.49.**Answer. $-4 - 2u^{-1} + 6u^{-2}$ **3.2.6.51.**Answer. $4x^{-2}(x^4 + 4)$ **3.2.6.53.**Answer. $a^{-3}(3 - 3a^4 + a^6)$ **3.2.6.55.**

Answer.

(a) No, because $\frac{1}{(x+y)^2}$ is not $\frac{1}{x^2} + \frac{1}{y^2}$.(b) Let $x = 1$, $y = 2$, then $(x + y)^{-2} = (1 + 2)^{-2} = 3^{-2} = \frac{1}{9}$, but $x^{-2} + y^{-2} = 1^{-2} + 2^{-2} = 1 + \frac{1}{4} = \frac{5}{4}$ **3.2.6.57.**

Answer.

(a) $x + x^{-1} = x + \frac{1}{x} = \frac{x^2}{x} + \frac{1}{x} = \frac{x^2 + 1}{x}$

(b) $x^3 + x^{-3} = x^3 + \frac{1}{x^3} = \frac{x^6}{x^3} + \frac{1}{x^3} = \frac{x^6 + 1}{x^3}$

(c) $x^n + x^{-n} = x^n + \frac{1}{x^n} = \frac{x^{2n}}{x^n} + \frac{1}{x^n} = \frac{x^{2n} + 1}{x^n}$

3.2.6.59.

Answer.

$$\begin{aligned} a^{-2}a^{-3} &= \frac{1}{a^2} \cdot \frac{1}{a^3} = \frac{1}{a^2 \cdot a^3} \\ &= \frac{1}{a^{2+3}} \\ &= \frac{1}{a^5} = a^{-5} \end{aligned}$$

3.2.6.61.

Answer.

$$\begin{aligned} \frac{a^{-2}}{a^{-6}} &= a^{-2} \div a^{-6} = \frac{1}{a^2} \div \frac{1}{a^6} \\ &\text{By the first law of exponents,} \\ &= \frac{1}{a^2} \cdot \frac{a^6}{1} = a^{6-2} \\ &= a^4 \end{aligned}$$

By the second law of exponents

3.3 · Roots and Radicals**3.3.10 · Roots and Radicals (Homework 3.3)****3.3.10.1.**

Answer.

(a) 11

(b) 3

(c) 5

3.3.10.3.

Answer.

(a) 2

(b) 2

(c) 9

3.3.10.5.

Answer.

(a) 3

(b) 3

(c) 2

3.3.10.7.

Answer.

(a) 2

(b) $\frac{1}{2}$

(c) $\frac{1}{8}$

3.3.10.9.

Answer.

(a) $\sqrt{3}$

(b) $4\sqrt[3]{x}$

(c) $\sqrt[5]{4x}$

3.3.10.11.

Answer.

(a) $\frac{1}{\sqrt[3]{6}}$

(b) $\frac{3}{\sqrt[8]{xy}}$

(c) $\sqrt[4]{x-2}$

3.3.10.13.

Answer.

(a) $7^{1/2}$

(b) $(2x)^{1/3}$

(c) $2z^{1/5}$

3.3.10.15.

Answer.

(a) $-3 \cdot 6^{-1/4}$

(b) $(x-3y)^{1/4}$

(c) $-(1+3b)^{-1/5}$

3.3.10.17.

Answer.

(a) 125

(b) 2

(c) 63

(d) $-2x^7$

3.3.10.19.

Answer.

(a) 1.414

(b) 4.217

(c) 1.125

(d) 0.140

(e) 2.782

3.3.10.21.

Answer.

(a) $g(x) = 3.7x^{1/3}$

(b) $\frac{H(x)}{85^{1/4}x^{1/4}}$


= (c) $F(t) = 25t^{-1/5}$

3.3.10.23.Answer. $x = 91.125$ **3.3.10.27.**Answer. $x = \frac{19}{2}$ **3.3.10.31.**Answer. $L = \frac{gT^2}{4\pi^2}$ **3.3.10.35.**Answer. $v = \frac{4}{3}\pi r^3$ **3.3.10.25.**Answer. $x = 241$ **3.3.10.29.**Answer. $x = \pm\sqrt{30}$ **3.3.10.33.**Answer. $s = \pm\sqrt{t^2 - r^2}$ **3.3.10.37.**Answer. $p = \frac{8Lv f}{\pi R^4}$ **3.3.10.39.**

Answer.

(a) $T = \frac{2\pi}{\sqrt{32}}L^{1/2}$

(b) 90 feet

**3.3.10.41.**


Answer.

(a) 3 meters per second

(d) 1.9 meters

(b) $b \approx 2.2$ meters per second

(e) 1.2 meters per second

**3.3.10.43.**


Answer.

(a) 6.5×10^{-13} cm; 1.17×10^{-36} cm³(b) 1.8×10^{14} g/cm³

(c)

Element	Carbon	Potassium	Cobalt	Technetium	Radium
Mass number, A	14	40	60	99	226
Radius, r (10^{-13} cm)	3.1	4.4	5.1	6	7.9

(d)




3.3.10.45.

Answer.

(a) $s = 50\sqrt{d}$

(b) 30 cm/sec




3.3.10.47.

Answer.

(a) $r = 12.1\sqrt{P}$

(b) 94 ft/sec




3.3.10.49.

Answer.

(a) 287; 343

(b) 2015; 2058

(c) The membership grows rapidly at first but is growing less rapidly with time.



3.3.10.51.

Answer.

(a) I

(b) III

(c) II

(d) none

3.3.10.53.

Answer.

(a) The graphs of $x^{1/n}$ become closer and closer to horizontal when n increases (for $x > 1$).

(b) 10, 4.64, 3.16, 2.51

(c) 1.58, 1.05, 1.005; the values decrease towards 1.

3.3.10.55.


Answer. The graphs of y_1 and y_2 are symmetric about $y_3 = x$.

3.3.10.57.

Answer. The graphs of y_1 and y_2 are symmetric about $y_3 = x$.

3.3.10.59.

Answer.



(b) $x = 36$

3.3.10.61.

Answer.

(a) $x^{1/2}$

(b) $(x^{1/2})^{1/2}$

(c)

$$\begin{aligned} \sqrt{\sqrt{x}} &= (x^{1/2})^{1/2} && \text{By definition of fractional exponents.} \\ &= x^{1/4} && \text{By the third law of exponents.} \\ &= \sqrt[4]{x} && \text{By definition of fractional exponents.} \end{aligned}$$

3.3.10.63.

Answer. $\frac{1}{4}x^{1/2} - 2x^{-1/2} + \frac{1}{\sqrt{2}}x$

3.3.10.65.

Answer. $3x^{-1/3} - \frac{1}{2}$

3.3.10.67.

Answer. $x^{0.5} + x^{-0.25} - x^0$

3.4 · Rational Exponents**3.4.8 · Rational Exponents (Homework 3.4)****3.4.8.1.**

Answer.

(a) 27

(b) 25

(c) 125

3.4.8.3.

Answer.

(a) $\frac{1}{64}$

(b) $\frac{1}{16}$

(c) $\frac{1}{256}$

3.4.8.5.

Answer.

(a) $\sqrt[5]{x^4}$

(b) $\frac{1}{\sqrt[6]{b^5}}$

(c) $\frac{1}{\sqrt[3]{(pq)^2}}$

3.4.8.7.

Answer.

(a) $3\sqrt[5]{x^2}$

(b) $\frac{4}{\sqrt[3]{z^4}}$

(c) $-2\sqrt[4]{xy^3}$

3.4.8.9.**Answer.**

(a) $x^{2/3}$

(b) $2a^{1/5}b^{3/5}$

(c) $-4mp^{-7/6}$

3.4.8.11.**Answer.**

(a) $(ab)^{2/3}$

(b) $8x^{-3/4}$

(c) $\frac{1}{3}RT^{-1/2}K^{-5/2}$

3.4.8.13.**Answer.**

(a) 8

(b) -81

(c) $2y^3$

3.4.8.15.**Answer.**

(a) $-a^4b^8$

(b) $2x^3y^9$

(c) $-3a^2b^3$

3.4.8.17.**Answer.**

(a) 7.931

(b) 10.903

(c) 0.090

(d) 35.142


3.4.8.19.**Answer.**

(a)

t	5	10	15	20
$I(t)$	131	199	254	302

Range: $[0, 302]$ (b) ≈ 19.812 or about 20 days

(c)

**3.4.8.21.****Answer.** All the graphs are increasing and concave up. For $x > 1$, each graph increases more quickly than the previous one.

3.4.8.23.**Answer.**

(a) $V = L^3, A = 6L^2$


(b) $L = V^{1/3}, L = \left(\frac{A}{6}\right)^{1/2}$

(c) $A = 6V^{2/3}$

(d) $\frac{A}{V} = \frac{6}{L}$. As L increases, the surface-to-volume ratio decreases.

3.4.8.25.**Answer.**

(a)




(b) \$7114.32

3.4.8.27.**Answer.**

(a)	A	10	100	1000	5000	10,000
	S	25	42	69	98	115

(b)



(c) 81, 71

(d) 126,000 sq km

3.4.8.29.**Answer.**

(a) Home range size: II, lung volume: III, brain mass: I, respiration rate: IV


(b) If $p > 1$, the graph is increasing and concave up. If $0 < p < 1$, the graph is increasing and concave down. If $p < 0$, the graph is decreasing and concave up.**3.4.8.31.****Answer.**

(a) Tricosanthes is the snake gourd and Lagenaria is the bottle gourd. Tricosanthes is thinner and Lagenaria is fatter.

- (b) $a \approx 9.5$
 (c) $a \approx 2$
 (d) Yes

3.4.8.33.**Answer.**

(a)


(b) ≈ 78.5 or about 79 species(c) 18.4°C

- (d) $f(9) \approx 85$, $f(10) \approx 79$, $f(19) \approx 49$, $f(20) \approx 47$; from 9°C to 10°C has the greater decrease, corresponding to the steeper slope. If the temperature range is 9°C , there will be approximately 85 species. If the temperature range is 10°C , there will be approximately 79 species. If the temperature range is 19°C , there will be approximately 49 species. If the temperature range is 20°C , there will be approximately 47 species.

3.4.8.35.**Answer.**

(a) $P = \frac{k}{\pi} d^{p-2}$

(b)



The power function is a good fit on this interval.

(c) 1.3

3.4.8.37.**Answer.**

(a) $4a^2$

(b) $9b^{5/3}$

3.4.8.39.**Answer.**

(a) $4w^{3/2}$

(b) $3z^2$

3.4.8.41.**Answer.**

(a) $\frac{1}{2k^{1/4}}$

(b) $\frac{4}{3h^{1/3}}$

3.4.8.43.**Answer.**

(a) Wren: 15 days, greylag goose: 28 days

(b) $\frac{I(m) \cdot W(m)}{m} = 0.18m^{-0.041}$

(c) Because $m^{-0.041}$ is close to m^0 , the fraction lost is close to 0.18.**3.4.8.45.****Answer.** $x = 64$ **3.4.8.47.****Answer.** $x = \frac{1}{243}$ **3.4.8.49.****Answer.** $x \approx 2.466$ **3.4.8.51.****Answer.**

(a) $p = 1.115 \times 10^{-12} a^{3/2}$

(b) 1.88 years

3.4.8.53.**Answer.** $\frac{13}{3}$ **3.4.8.55.****Answer.** 0.665**3.4.8.57.****Answer.** $2x^{3/2} - 2x$ **3.4.8.59.****Answer.** $\frac{1}{2}y^{1/3} + \frac{3}{2}y^{-7/6}$ **3.4.8.61.****Answer.** $2x^{1/2} - x^{1/4} - 1$ **3.4.8.63.****Answer.** $a^{3/2} - 4a^{3/4} + 4$ **3.4.8.65.****Answer.** $x(x^{1/2} + 1)$ **3.4.8.67.****Answer.** $\frac{y-1}{y^{1/4}}$ **3.4.8.69.****Answer.** $\frac{a^{2/3} + a^{1/3} - 1}{a^{1/3}}$

3.5 · Chapter Summary and Review

3.5.2 · Chapter 3 Review Problems

3.5.2.1.**Answer.**


a $d = 1.75t^2$

b 63 cm

3.5.2.3.**Answer.** 480 bottles**3.5.2.5.****Answer.**

a $w = \frac{k}{r^2}$

b



c $3960\sqrt{3} \approx 6860$ miles

3.5.2.7.

Answer.
 $y = 1.2x^2$

3.5.2.9.

Answer.
 $y = \frac{20}{x}$

3.5.2.11.

Answer.

a $\frac{1}{81}$

3.5.2.13.

Answer.

a $\frac{1}{243m^5}$

3.5.2.15.

Answer.

a $\frac{2}{c^3}$

b $\frac{1}{64}$

b $\frac{-7}{y^8}$

b $\frac{99}{z^2}$

3.5.2.17.

Answer.

a $25\sqrt{m}$

3.5.2.19.

Answer.

a $\frac{1}{\sqrt[4]{27q^3}}$

b $7\sqrt{u^3v^3}$

3.5.2.21.

Answer.

a $2x^{2/3}$

b $\frac{1}{4}x^{1/4}$

3.5.2.23.


Answer.

a $6b^{-3/4}$

b $\frac{-1}{3}b^{-1/3}$


3.5.2.25.

Answer.



3.5.2.27.

Answer.



3.5.2.29.

Answer. $f(x) = \frac{2}{3}x^{-4}$

3.5.2.31.

Answer.


(a)	x	16	$\frac{1}{4}$	3	(a)	$\frac{100}{f(x)}$
	$Q(x)$	4096	$\frac{1}{8}$	$4\sqrt{3^5} \approx 62.35$		400,000

3.5.2.33.


Answer.

0	1	5	10	20	50	70	100
0	1	1.62	2.00	2.46	3.23	3.58	3.99

(b)



(b)




3.5.2.35.

Answer. 112 kg

3.5.2.37.

Answer.

a

b 283.7 or ≈ 284

c 2051

3.5.2.39.**Answer.**

a It is the cost of producing the first ship.

b $C = \frac{12}{\sqrt[8]{x}}$ million


c About \$11 million; about 8.3%

d About 8.3%

3.5.2.41.**Answer.** $t = 10$ **3.5.2.45.****Answer.** $x = 5$ **3.5.2.49.****Answer.** $y = 29,524$ **3.5.2.51.****Answer.** $g = \frac{2v}{t^2}$ **3.5.2.55.****Answer.** $49t^2$ **3.5.2.57.****Answer.** $\frac{k^7}{64}$ **3.5.2.59.****Answer.** $8a^2$ **3.5.2.43.****Answer.** $x = 7$ **3.5.2.47.****Answer.** $x = 75$ **3.5.2.53.****Answer.** $p = \pm 2\sqrt{R^2 - R}$ **3.5.2.61.****Answer.**

(a) 132.6 km

(b)

**3.5.2.63.****Answer.**

(a) 480

(b) 498

3.5.2.65.**Answer.**

a \$450

b $t = 8$: It costs \$864 to insulate a ceiling with 8 cm of insulation over an area of 600 square meters.c $C = 0.72A$ d $C = 18T$ e $C = 0.18AT$

f \$1440

3.5.2.67.**Answer.**a $N = \frac{k}{d^2 E^3}$, where N is number of people, d is distance in miles from the road, E is the elevation gain, and k is the constant of variation.b $k \approx 0.01$

c 3


4 · Exponential Functions**4.1 · Exponential Growth and Decay****4.1.8 · Exponential Growth and Decay (Homework 4.1)****4.1.8.1.****Answer.**

(a) \$28

(b) \$31.36

4.1.8.3.**Answer.** It is 99% of what it was 2 years ago.**4.1.8.5.****Answer.**(a) $P = 1200 + 150t$; 1650(b) $P = 1200 \cdot 1.5^t$; 4050**4.1.8.7.****Answer.**(a) $V = 18,000 - 2000t$; \$8000(b) $V = 18,000 \cdot 0.8^t$; \$5898.24**4.1.8.9.****Answer.** A: 20%; B: 2%; C: 7.5%; D: 100%; E: 115%**4.1.8.11.****Answer.**

a $P = 20,000 \cdot 2.5^{t/6}$




c 36,840 bees; 424,128 bees

4.1.8.13.

Answer.

a $A = 4000 \cdot 1.08^t$

c \$4665.60; \$8635.70




4.1.8.15.

Answer.

a $P = 200,000 \cdot 1.05^t$


c \$359,171; \$746,691



4.1.8.17.

Answer.

a $P = 250,000 \cdot 0.75^{t/2}$




c 162,380; 79,102

4.1.8.19.

Answer.

a $L = 0.85^{d/4}$

c 44%; 16%




4.1.8.21.

Answer.

a $P = 50 \cdot 0.992^t$

c 46.1 lb; 22.4 lb

**4.1.8.23.****Answer.**

a 3^{x+4}

b 3^{4x}

c 12^x

4.1.8.25.**Answer.**

a b^{-2t}

b $b^{t/2}$

c 1

4.1.8.27.

Answer. $P(t+1) = 12(3)^{t+1} = 12(3)^t \cdot 3 = P(t) \cdot 3$

4.1.8.29.

Answer. $P(x+k) = P_0 a^{x+k} = P_0 a^x \cdot a^k = P(x) \cdot a^k$

4.1.8.31.**Answer.**

- a In the expression $2 \cdot 3^t$, only the 3 is raised to a power t , and the result is doubled, but if both the 2 and the 3 were raised to the power t , the result would be 6^t .

b

t	0	1	2
$P(t)$	2	6	18
$Q(t)$	1	6	36

4.1.8.33.

Answer. 4

4.1.8.37.

Answer. $r \approx 0.14$

4.1.8.35.

Answer. 1.2

4.1.8.39.

Answer. $r \approx 0.04$

4.1.8.41.**Answer.**

a $P(t) = 1,545,387b^t$

b Growth factor 1.049; Percent rate of growth 4.9%

c 2,493,401

4.1.8.43.**Answer.**

a 365

b $N(t) = 365(0.356)^t$

c

4.1.8.45.

Answer. The growth factor is 1.5.

t	0	1	2	3	4
P	8	12	18	27	40.5

4.1.8.49.

Answer. The decay factor is 0.8.

w	0	1	2	3	4
N	120	96	76.8	61.44	49.15

4.1.8.53.

Answer. The growth factor is 1.1.

n	0	1	2	3	4
B	200	220	242	266.2	292.82

4.1.8.55.

Answer.

(a) Initial value 4, growth factor $2^{1/3}$

(b) $f(x) = 4 \cdot 2^{x/3}$

4.1.8.59.

Answer. 84.6%, 55.8%

4.1.8.61.

Answer. No, an increase of 48% in 6 years corresponds to a growth factor of $1.48^{1/6} \approx 1.0675$, or an annual growth rate of about 6.75%.

4.1.8.63.

Answer.

a $P(t) = 16,986,335(1+r)^t$

4.1.8.65.

Answer.

a 3.53%

b 3.53%

c No

d 3.53%


4.1.8.67.

Answer.

a 39; 1.045

b 35; 1.047

c Species B



d 0.03. (Therefore, none)

4.1.8.47.

Answer. The growth factor is 1.2.

x	0	1	2	3	4
Q	20	24	28.8	34.56	41.47

4.1.8.51.

Answer. The decay factor is 0.8.

t	0	1	2	3	4
C	10	8	6.4	5.12	4.10

4.1.8.57.

Answer.

(a) Initial value 80, decay factor $\frac{1}{2}$


(b) $f(x) = 80 \cdot \left(\frac{1}{2}\right)^x$

4.1.8.69.**Answer.**

a

t	0	2	4	6	8
$L(t)$	3	6	9	12	15

$$L(t) = 3 + 1.5t$$



b

t	0	2	4	6	8
$E(t)$	3	6	12	24	48

$$E(t) = 3 \cdot 2^{t/2}$$

4.1.8.71.**Answer.**

a 244 tigers per year

c Linear: 3067; Exponential:
4170

b 0.97; 3%

4.2 · Exponential Functions**4.2.6 · Exponential Functions (Homework 4.2)****4.2.6.1.****Answer.**


(a) 26; increasing

(c) 75; decreasing

(b) 1.2; decreasing

(d) $\frac{2}{3}$; increasing**4.2.6.3.****Answer.**

x	-3	-2	-1	0	1	2	3
$f(x) = 3^x$	$\frac{1}{27}$	$\frac{1}{9}$	$\frac{1}{3}$	1	3	9	27
$g(x) = \left(\frac{1}{3}\right)^x$	27	9	3	1	$\frac{1}{3}$	$\frac{1}{9}$	$\frac{1}{27}$




The two graphs are reflections of each other across the y -axis. f is increasing, g is decreasing. f has the negative x -axis as an asymptote, and g has the positive x -axis as its asymptote.

4.2.6.5.

Answer.

t	-3	-2	-1	0	1	2	3
$h(t) = 4^{-t}$	64	16	4	1	$\frac{1}{4}$	$\frac{1}{16}$	$\frac{1}{64}$
$q(t) = -4^t$	$-\frac{1}{64}$	$-\frac{1}{16}$	$-\frac{1}{4}$	-1	-4	-16	-64



The graphs are reflections of each other across the origin. Both are decreasing, but h has the negative t -axis as an asymptote, and q has the positive t -axis as its asymptote.

4.2.6.7.

Answer.

a I

b IV


c III

d II

4.2.6.9.

Answer.

a




b [1.08, 14.85]

4.2.6.11.

Answer.

a



b [16.38, 152.59]

4.2.6.13.

Answer. Because they are defined by equivalent expressions, (b), (c), and (d) have identical graphs


4.2.6.15.

Answer.

a To evaluate f we subtract 1 from the input before evaluating the exponential function; to evaluate g we subtract 1 from the output of the exponential function.

b

x	$y = 2^x$	$f(x)$	$g(x)$
-2	$\frac{1}{4}$	$\frac{1}{8}$	$-\frac{3}{4}$
-1	$\frac{1}{2}$	$\frac{1}{4}$	$-\frac{1}{2}$
0	1	$\frac{1}{2}$	0
1	2	1	1
2	4	2	3



c The graph of f is translated 1 unit to the right; the graph of g is shifted 1 unit down.


4.2.6.17.

Answer.

a To evaluate f we take the negative of the output of the exponential function; to evaluate g we take the negative of the input.

x	$y = 3^x$	$f(x)$	$g(x)$
-2	$\frac{1}{9}$	$-\frac{1}{9}$	9
-1	$\frac{1}{3}$	$-\frac{1}{3}$	3
0	1	-1	1
1	3	-3	$\frac{1}{3}$
2	9	-9	$\frac{1}{9}$

b



c The graph of f is reflected about the x -axis; the graph of g is reflected about the y -axis.

4.2.6.19.

Answer.

a $3(5^{a+2})$ is not equivalent to $9 \cdot 3(5^a)$.

b $3(5^{2a})$ is not equivalent to $2 \cdot 3(5^a)$.

4.2.6.21.

Answer.

a $8^w - 8^z$ is not equivalent to 8^{w-z} .

b 8^{-x} is equivalent to $\frac{1}{8^x}$.

4.2.6.23.**Answer.**

a $P_0 = 300$

c $b = 2$

b

x	0	1	2
$f(x)$	300	600	1200

d $f(x) = 300(2)^x$

4.2.6.25.**Answer.**

a $S_0 = 150$

b $b \approx 0.55$

c $S(d) = 150(0.55)^d$

4.2.6.27.

Answer. $\frac{2}{3}$

4.2.6.29.

Answer. $-\frac{1}{4}$

4.2.6.31.

Answer. $\frac{1}{7}$

4.2.6.33.


Answer. $-\frac{5}{4}$

4.2.6.35.

Answer. ± 2

4.2.6.37.**Answer.**


a $N(t) = 26(2)^{t/6}$



c 72 days later

4.2.6.39.**Answer.**

a $V(t) = 700(0.7)^{t/2}$



c 4 yr

4.2.6.41.

Answer.
 $x = 2.26$

4.2.6.43.

Answer.
 $x = -1.40$

4.2.6.45.**Answer.**

a Power

c Power

b Exponential

d Neither

4.2.6.47.**Answer.**


- a Exponential $y = 3 \cdot 2^x$
 b Power $P = 0.5t^2$

4.2.6.49.**Answer.**

- a Power $y = 100x^{-1}$
 b Exponential $P = \frac{1}{4} \cdot 2^x$

4.2.6.51.**Answer.**

x	$f(x) = x^2$	$g(x) = 2^x$
-2	4	$\frac{1}{4}$
-1	1	$\frac{1}{2}$
0	0	1
1	1	2
2	4	4
3	9	8
4	16	16
5	25	32




- a Range of f : $[0, \infty)$; Range of g : $(0, \infty)$
 b 3
 c $-0.7667, 2, 4$
 d $(-0.7667, 2)$ and $(4, \infty)$
 e g

4.2.6.53.


Answer.

a $y = 3^x - 4$




Domain: $(-\infty, \infty)$; range: $(-4, \infty)$, x -intercept $(1.26, 0)$; y -intercept $(0, -3)$; horizontal asymptote $y = -4$

b $y = 3^{x-4}$,



Domain: $(-\infty, \infty)$; range: $(0, \infty)$, no x -intercept; y -intercept $(0, \frac{1}{81})$; the x -axis is the horizontal asymptote.

c $y = -4 \cdot 3^x$,




Domain: $(-\infty, \infty)$; range: $(-\infty, 0)$, no x -intercept; y -intercept $(0, -4)$; the x -axis is the horizontal asymptote.

4.2.6.55.


Answer.

a $y = -6^t$




Domain: $(-\infty, \infty)$; range: $(-\infty, 0)$, no t -intercept; y -intercept $(0, -1)$; the t -axis is the horizontal asymptote.

b $y = 6^{-t}$,



Domain: $(-\infty, \infty)$; range: $(0, \infty)$, no t -intercept; y -intercept $(0, 1)$; the t -axis is the horizontal asymptote.


c $y = -6^{-t}$,



Domain: $(-\infty, \infty)$; range: $(-\infty, 0)$, no t -intercept; y -intercept $(0, -1)$; the t -axis is the horizontal asymptote.


4.2.6.57.**Answer.**

a $y = 2^{x-3}$



Domain: $(-\infty, \infty)$; range: $(0, \infty)$, no x -intercept; y -intercept $(0, \frac{1}{8})$; the x -axis is the horizontal asymptote.


b $y = 2^{x-3} + 4$,



Domain: $(-\infty, \infty)$; range: $(4, \infty)$, no x -intercept; y -intercept $> (0, \frac{33}{8})$; horizontal asymptote $y = 4$


4.2.6.59.**Answer.**

a $y = -\left(\frac{1}{2}\right)^t$



Domain: $(-\infty, \infty)$; range: $(-\infty, 0)$, no t -intercept; y -intercept $(0, -1)$; the t -axis is the horizontal asymptote.

b $y = 6 - \left(\frac{1}{2}\right)^t$,



Domain: $(-\infty, \infty)$; range: $(-\infty, 6)$, t -intercept approximately $(-2.58, 0)$; y -intercept $(0, 5)$; horizontal asymptote is $y = 6$

4.2.6.61.**Answer.**

a The graph of $y = 2^x$ has been reflected about the y -axis and shifted up 2 units.

b $y = 2^{-x} + 2$

4.2.6.63.**Answer.**

a The graph of $y = 2^x$ has been reflected about the x -axis and shifted up 10 units.

b $y = -2^x + 10$

4.2.6.65.**Answer.**

a I

b III


c II

4.2.6.67.**Answer.**

a

t	3.5	4	8	10	15
$f(t)$	128	154.75	184.05	150.93	103.96

b



c From 0 to 3 minutes, the volunteer is walking with heart rate 100 beats per minute. The volunteer jogged at a steady pace from 3 to 4 minutes, and the heart rate increased to about 155 beats per minutes. From 4 to 9 minutes, the jogging pace increased, and the heart rate rose to about 185 beats per minute. The cooldown started at 9 minutes, and the heart rate decreased rapidly and leveled off to about 100 beats per minute.

4.3 · Logarithms

4.3.7 · Logarithms (Homework 4.3)

4.3.7.1.

Answer.

(a) 2

(b) 5

4.3.7.5.

Answer.

(a) 1

(b) 0

4.3.7.9.

Answer.

(a) -1

(b) -3

4.3.7.11.

Answer.

$$\log_2(1024) = 10$$

4.3.7.17.

Answer.

$$\log_{0.8}(M) = 1.2$$

4.3.7.23.

Answer.

$$\begin{array}{cc} \text{a} & \text{b} \\ \log_4(2.5) & 0.7 \end{array}$$

4.3.7.3.

Answer.

(a) $\frac{1}{2}$

(b) -1

4.3.7.7.

Answer.

(a) 5

(b) 6

4.3.7.13.

Answer.

$$\log(5) \approx 0.699$$

4.3.7.19.

Answer.

$$\log_x(W - 3) = 5t$$

4.3.7.25.

Answer.

$$\begin{array}{cc} \text{a} & \text{b} \\ \log(0.003) & -2.5 \end{array}$$

4.3.7.15.

Answer.

$$\log_t(16) = \frac{3}{2}$$

4.3.7.21.

Answer.

$$\log_3(2N_0) = -0.2t$$

4.3.7.27.**Answer.**

$$\text{a } 0 < \frac{\log(7)}{1} <$$

$$\text{b } 0.85$$

4.3.7.29.**Answer.**

$$\text{a } 3 < \frac{\log_3(67.9)}{4} <$$

$$\text{b } 3.84$$

4.3.7.31.**Answer.**

$$\text{(a) } 0.7348$$

$$\text{(b) } 1.7348$$

$$\text{(c) } 2.7348$$

$$\text{(d) } 3.7348$$

When the input to the common logarithm is multiplied by 10, the output is increased by 1.

4.3.7.35.**Answer.** -0.23 **4.3.7.39.****Answer.** 0.77 **4.3.7.43.****Answer.** 3.63 **4.3.7.45.**

Answer. $2 \cdot 5^x \neq 10^x$; the first step should be to divide both sides of the equation by 2; $x = \log_5(424)$.

4.3.7.33.**Answer.**

$$\text{(a) } 0.3010$$

$$\text{(b) } 0.6021$$

$$\text{(c) } 0.9031$$

$$\text{(d) } 1.2041$$

When the input to the common logarithm is doubled, the output is increased by about 0.3010.

4.3.7.37.**Answer.** 2.53 **4.3.7.41.****Answer.** -0.68 **4.3.7.47.**

Answer. $\frac{10^{4x}}{4} \neq 10^x$; the first step should be to write $4x = \log(20)$; $x = \frac{\log(20)}{4}$.

4.3.7.49.**Answer.**


$$\text{(a) } 33,855,812$$

$$\text{(b) } 38,515,295; 41,080,265; 43,816,051$$

$$\text{(c) } 2002$$

$$\text{(d) } 2012$$


(e)

**4.3.7.51.****Answer.**

(a) 85.5

(c) 1.45

(b) Decreasing; range: $[5.4, 1355.2]$



(d) $\frac{1}{100}$

(e) $10^{0.4} \approx 2.5119$

(f) 2.15×10^{-6} to 855,067

4.3.7.53.

Answer. 9.60 in

4.3.7.55.

Answer. 1.91 mi

4.3.7.57.

Answer. 3.34 mi

4.3.7.59.

Answer. 1

4.3.7.61.

Answer. 0

4.3.7.63.

Answer. 1

4.3.7.65.

Answer. 0

4.4 · Properties of Logarithms

4.4.7 · Properties of Logarithms (Homework 4.4)

4.4.7.1.

Answer.

(a) 10^8

(b) 2; 6; 8; $2 + 6 = 8$

4.4.7.3.

Answer.

(a) b^3

(b) 8; 5; 3; $8 - 5 = 3$

4.4.7.5.

Answer.

(a) 10^{15}

(b) 15; 3; $15 = 3 \cdot 5$

4.4.7.7.

Answer.

4.4.7.9.

Answer.

(a) $\log_b(2) + \log_b(x)$

(b) $\log_b(2) - \log_b(x)$

(a) $1 + 4 \log_3(x)$

(b) $\frac{1}{t} \log_5(1.1)$

4.4.7.11.

Answer.

4.4.7.13.

Answer.

(a) $\frac{1}{2} + \frac{1}{2} \log_b(x)$

(b) $\frac{1}{3} \log_3((x^2 + 1))$

(a) $\log(P_0) + t \log(1 - m)$

(b) $4t[\log_4(4 + r) - 1]$

4.4.7.15.**Answer.**

(a) $\log_b(4)$ (b) $\log_4(x^2y^3)$ (a) $\log(2x^{5/2})$ (b) $\log(t-4)$

4.4.7.19.**Answer.**

(a) $\log\left(\frac{1}{27}\right)$ (b) $\log_6(2w^2)$

4.4.7.21.**Answer.**


(a) 1.7917 (b) -0.9163 (a) 2.1972 (b) 1.9560

4.4.7.25.**Answer.** 2.8074**4.4.7.27.****Answer.** 0.8928**4.4.7.29.****Answer.** ± 1.3977 **4.4.7.31.****Answer.** -1.6092**4.4.7.33.****Answer.** 0.2736**4.4.7.35.****Answer.** -12.4864**4.4.7.37.****Answer.**

(a) $S(t) = S_0(1.09)^t$ (b) 4.7 hours

4.4.7.39.**Answer.**

(a) $C(t) = 0.7(0.80)^t$



(b) After 2.5 hours

4.4.7.41.**Answer.**

(a) $J(t) = 1,041,000 \cdot 1.0182^t$ (b) In 2040

4.4.7.43.**Answer.**

(a) $S(t) = S_0 \cdot 0.9527^t$ (b) 28.61 hours

4.4.7.45.**Answer.**

(a) 5 (b) 6 (c) 5

(a) and (c) are equal.

4.4.7.47.**Answer.**

(a) 6 (b) 9 (c) 6

(a) and (c) are equal.

4.4.7.49.**Answer.**

(a) $\log(24) \approx 1.38$ (b) $\log(240) \approx 2.38$ (c) $\log(230) \approx 2.36$

None are equal.

4.4.7.51.**Answer.**

(a) $\log(60) \approx 1.78$ (b) $\log(5) \approx 0.70$ (c) $\frac{\log(75)}{\log(15)} \approx 1.59$

None are equal.

4.4.7.53.**Answer.** 12.9%**4.4.7.55.****Answer.** About 11 years**4.4.7.57.****Answer.**

(a) $A = 1000 \left(1 + \frac{0.12}{n}\right)^{5n}$

 A increases.

(b)

X	Ψ_1
1	1762.3
2	1790.8
4	1806.1
8	1816.7
12	1820.8
16	1821.9

(c) 16; 31; 553

(d) Increasing, concave down, asymptotically approaching $A \approx 1822.12$ **4.4.7.59.**

Answer. $k = \frac{1}{t} \frac{\log(N/N_0)}{\log(a)}$

4.4.7.61.

Answer. $t = \frac{1}{k} \log\left(\frac{A}{A_0} + 1\right)$

4.4.7.63.

Answer. $q = \frac{\log(w/p)}{\log(v)}$

4.4.7.65.**Answer.**

(a) $x = b^m, y = b^n$

(d) $\log_b(b^{m+n}) = m + n$

(b) $\log_b(b^m \cdot b^n)$

(c) $\log_b(b^m \cdot b^n) = \log_b(b^{m+n})$

(e) $\log_b(b^{m+n}) = \log_b(x) + \log_b(y)$

4.4.7.67.**Answer.**

(a) $x = b^m$

(d) $\log_b(b^{mk}) = mk$

(b) $\log_b(b^m)^k$

(c) $\log_b(b^m)^k = \log_b(b^{mk})$

(e) $\log_b(b^{mk}) = (\log_b(x)) \cdot k$

4.5 · Exponential Models

4.5.6 · Exponential Models (Homework 4.5)

4.5.6.1.

Answer. $A(x) = 0.14(50)^{x/3}$

4.5.6.5.

Answer. $M(x) = 62,500(0.2)^x$

4.5.6.9.


Answer. $y = \frac{4}{3}(3)^{x/4}$

4.5.6.13.

Answer.

(a) $y = 2.6 - 1.3x$

(b) $y = 2.6(0.5)^x$



4.5.6.3.

Answer. $f(x) = \frac{65,536}{729} \left(\frac{3}{4}\right)^x$

4.5.6.7.

Answer. $s(x) = \frac{1}{135}(9)^x$

4.5.6.11.


Answer. $y = 50(2)^{-x/4}$

4.5.6.15.

Answer.

(a) $y = -36 - 16x$

(b) $y = \frac{12}{5}(5)^{-x/3}$




4.5.6.17.

Answer.

(a) $y = 2.5 + 0.875x$

(b) $y = 1.5(2)^{x/2}$




4.5.6.19.

Answer.

(a) $P = P_0(1.052)^t$; t is the number of years since 1990.

(b) $\frac{\log(2)}{\log(1.052)} \approx 13.7$ years



4.5.6.21.


Answer.

(a) $GDP = 1.028^t$ million pounds

(b) $\frac{\log(2)}{\log(1.028)} \approx 25.1$ years

(c) 50.2 years

(d)




4.5.6.23.

Answer.

a $\frac{\log(0.5)}{\log(0.946)} \approx 12.5$ hours

b 25 hours

c




4.5.6.25.

Answer.

a $\frac{\log(0.5)}{\log(0.844)} \approx 4.1$ hours

b 8.2 hours

c



4.5.6.27.

Answer.

a $P = 2000(2)^{t/5}$

b 14.87%

4.5.6.29.

Answer.

a $D = D_0 \left(\frac{1}{2}\right)^{t/18}$

b 3.78%

4.5.6.31.

Answer.

(a) $A = A_0 \left(\frac{1}{2}\right)^{t/1620}$

(b) 0.043%

4.5.6.33.

Answer.

(a) $P = P_0(2)^{t/25}$

(b) 2.81%

4.5.6.35.**Answer.**

(a) $ab^D = 2 \cdot ab^0 = 2a$

(b) $b^D = 2$

(c) $f(t+D) = ab^{t+D} = a \cdot b^t \cdot b^D = ab^t \cdot 2 = 2f(t)$

(d) For any value of t , after D units of time, the new value of f is 2 times the old value.**4.5.6.37.****Answer.**

(a) $ab^R = \frac{1}{3} \cdot ab^0 = \frac{1}{3}a$

(b) $b^R = \frac{1}{3}$

(c) $g(t+R) = ab^{t+R} = a \cdot b^t \cdot b^R = ab^t \cdot \frac{1}{3} = \frac{1}{3}g(t)$

(d) For any value of t , after R units of time, the new value of g is $\frac{1}{3}$ times the old value.**4.5.6.39.****Answer.**

(a) $A = A_0 \left(\frac{1}{2}\right)^{t/5730}$

(b) About 760 years old

4.5.6.41.**Answer.**

(a) $A = A_0 \left(\frac{1}{2}\right)^{t/432}$

(b) About 220 years

4.5.6.43.**Answer.** ≈ 30 years; ≈ 33 years**4.5.6.45.****Answer.** \$445.89; \$376.50**4.5.6.47.****Answer.**

(a) $N(t) = 2200(2)^{t/1.5}$

(b) The given model has a smaller growth factor, 1.356, than $2^{1/1.5} \approx 1.59$.

	Name of chip	Year	Moore's law	$N(t)$	Actual number
(c)	Pentium IV	2000	2,306,867,200	20,427,413	42,000,000
	Pentium M (Banias)	2003	9,227,468,800	50,932,200	77,000,000
	Pentium M (Dothan)	2004	14,647,693,680	69,064,063	140,000,000

(d) About 2.3 years

4.6 · Chapter Summary and Review

4.6.2 · Chapter 4 Review Problems

4.6.2.1.

Answer.

a $D = 8(1.5)^{t/5}$

b 18; 44

4.6.2.3.

Answer.

a $M = 100(0.85)^t$

b 52.2 mg; 19.7 mg

4.6.2.5.

Answer. $16n^{2x+10}$

4.6.2.7.

Answer. $\frac{1}{m^{x+2}}$

4.6.2.9.

Answer. $g(t) = 16(0.85)^t$

4.6.2.11.

Answer. $f(x) = 500 \left(\frac{1}{5}\right)^x$

4.6.2.13.


Answer. 4.8% loss

4.6.2.15.

Answer. 6% loss

4.6.2.17.

Answer.




(b) y -intercept $(0, 6)$; asymptote:
 $y = 0$

(c) $[3.472, 10.368]$

4.6.2.19.

Answer.



(b) x -intercept $\left(\frac{\log(3)}{\log(2)}, 0\right)$;
 y -intercept $(0, -2)$;
asymptote: $y = -3$

(c) $[-2.875, 5]$

4.6.2.21.

Answer. $\frac{-4}{3}$


4.6.2.23.

Answer. -11

4.6.2.25.

Answer.


(a)



Not (quite) equivalent

(b) $2^{1/8} \approx 1.090507733 > 1.0905$ **4.6.2.27.****Answer.**

(a)




Equivalent



$$(b) \left(\frac{1}{3}\right)^{x-2} = \left(\frac{1}{3}\right)^x \cdot \left(\frac{1}{3}\right)^{-2} = \left(\frac{1}{3}\right)^x \cdot 9$$

4.6.2.29.**Answer.**

(a) $y = 4 + 2^{x+1}$

(b) Shift the graph of f 1 unit left, 4 units up.**4.6.2.31.****Answer.**

(a) $y = 6 - 3 \cdot 2^x$

(b) Scale vertically by 3, reflect about x -axis, shift 6 units up.**4.6.2.33.****Answer.** g eventually grows faster.**4.6.2.35.****Answer.** $2^{1.5} \approx 2.83$; 2.25

4.6.2.37.

Answer. $M = M_0(2)^{t/10}$, where M is the organic content, M_0 is the organic content at 0°C , and t is the temperature in $^\circ\text{Celsius}$.

4.6.2.39.**Answer.** 4**4.6.2.41.****Answer.** -1**4.6.2.43.****Answer.** -3**4.6.2.45.****Answer.** $\log_{0.3}(x+1) = -2$ **4.6.2.47.****Answer.** $\frac{\log(5.1)}{1.3} \approx 0.5433$ **4.6.2.49.****Answer.** $\frac{\log(2.9/3)}{-0.7} \approx 0.21$ **4.6.2.51.****Answer.** $\log_b(x) + \frac{1}{3}\log_b(y) - 2\log_b(z)$ **4.6.2.53.****Answer.** $\frac{4}{3}\log(x) - \frac{1}{3}\log(y)$ **4.6.2.55.****Answer.** $\log\left(\sqrt[3]{\frac{x}{y^2}}\right)$ **4.6.2.57.****Answer.** $\log\left(\frac{1}{8}\right)$ **4.6.2.59.****Answer.** $\frac{\log(63)}{\log(3)} \approx 3.77$ **4.6.2.61.****Answer.** $\frac{\log(50)}{-0.3\log(6)} \approx -7.278$ **4.6.2.63.****Answer.** $\frac{\log(N/N_0)}{k}$ **4.6.2.65.****Answer.**

(a) 238

(b) 2010

4.6.2.67.**Answer.**(a) $C = 90(1.06)^t$

(b) \$94.48

(c) 5 years

4.6.2.69.**Answer.**

(a) 7.4 years

(b) 6.1

4.6.2.71.**Answer.** $f(x) \approx 1600(1.035)^x$ **4.6.2.73.****Answer.** $g(x) \approx 600(0.075)^x$ **4.6.2.75.****Answer.**(a) $\frac{\log(2)}{\log(1.001)} \approx 693$ years

(b) 105 years

4.6.2.77.**Answer.** 17%

4.6.2.79.**Answer.** \$2192.78**4.6.2.81.****Answer.**

(a)

Day	1	2	3	...	t	...	30
Wage (cent)	2	4	8	...	2^t	...	2^{30}

(b) $W(t) = 2^t$ cents

(c) \$327.68; \$10,737,418.24

5 · Logarithmic Functions**5.1 · Inverse Functions****5.1.10 · Inverse Functions (Homework 5.1)****5.1.10.1.****Answer.**

(a)

x	-1	0	1	2
$f(x)$	0	1	-2	-1

y	0	1	-2	-1
$f^{-1}(y)$	-1	0	1	2

(b) $f^{-1}(1) = 0$ (c) $f^{-1}(-1) = 2$ **5.1.10.3.****Answer.**

(a)

x	-1	0	1	2
$f(x)$	-1	1	3	11

y	-1	1	3	11
$f^{-1}(y)$	-1	0	1	2

(b) $f^{-1}(1) = 0$ (c) $f^{-1}(3) = 1$ **5.1.10.5.****Answer.**(a) $f(60) \approx 38$. The car that left the 60-foot skid marks was traveling at 38 mph.(b) $f^{-1}(60) \approx 150$. The car traveling at 60 mph left 150-foot skid marks**5.1.10.7.****Answer.**

(a) (60 hours, 78 grams)

(b) $f^{-1}(90) \approx 19$, so that the vampire bat's weight has dropped to 90 grams about 19 hours after its last meal.**5.1.10.9.****Answer.**

- (a) $g(0.05) = 0.28$. At 5% interest, \$1 earns \$0.28 interest in 5 years.
- (b) 8.45%
- (c) $g^{-1}(I) = (I + 1)^{1/5} - 1$
- (d) $g^{-1}(0.50) \approx 0.0845$

5.1.10.11.**Answer.**

- (a) $f(0.5) \approx 62.9$. At an altitude of 0.5 miles, you can see 62.9 miles to the horizon.
- (b) 0.0126 mile, or 66.7 feet
- (c) $h = f^{-1}(d) = \frac{d^2}{7920}$
- (d) $f^{-1}(10) \approx 0.0126$


5.1.10.13.**Answer.**

- (a) $h^{-1}(3) \approx -4$
- (b) $h^{-1}(x) = 5 - x^2$; $h^{-1}(3) = -4$

5.1.10.15.**Answer.**

- (a) $f^{-1}(y) = 3\sqrt[3]{y} + 2$
- (b) $f^{-1}(f(4)) = f^{-1}(8) = 4$
- (c) $f(f^{-1}(-8)) = f(0) = -8$

(d)

**5.1.10.17.****Answer.** 6**5.1.10.19.****Answer.** $\frac{2}{9}$ **5.1.10.21.****Answer.** 4


5.1.10.23.**Answer.**

(a)

x	0	6
y	300	1200

(b)

x	300	1200
y	0	6


**5.1.10.25.****Answer.**

(a)


x	0	1	2
y	5	20	100

(b)


x	5	20	100
y	0	1	2

**5.1.10.27.****Answer.**


(a) $f^{-1}(x) = \frac{x+6}{2}$

**5.1.10.29.****Answer.**

(a) $f^{-1}(x) = \sqrt[3]{x-1}$

**5.1.10.31.****Answer.**


(a) $f^{-1}(x) = \frac{1}{x} + 1$

**5.1.10.33.****Answer.**

(a) Domain: $(-\infty, 4]$; Range: $[0, \infty)$

(b) $g^{-1}(x) = 4 - x^2$

(c) Domain: $[0, \infty)$; Range: $(-\infty, 4]$



5.1.10.35.

Answer. (a) and (d)

5.1.10.39.

Answer. (a)

5.1.10.37.

Answer. (a)

5.1.10.41.

Answer. (a) and (b)

5.1.10.43.

Answer.

(a) $f(x) = 4 + 2x$; IV

(c) $f(x) = -4 - 2x$; I

(b) $f(x) = 2 - \frac{x}{2}$; III

(d) $f(x) = \frac{x}{2}$; II

5.1.10.45.

Answer.

(a) III

(b) II

(c) I

5.2 · Logarithmic Functions


5.2.7 · Logarithmic Functions (Homework 5.2)

5.2.7.1.**Answer.**

x	-1	0	1	2
2^x	$\frac{1}{2}$	1	2	4

x	$\frac{1}{2}$	1	2	4
$\log_2(x)$	-1	0	1	2


(b)

**5.2.7.3.****Answer.**

x	-2	-1	0	1
$(\frac{1}{3})^x$	9	3	1	$\frac{1}{3}$

x	9	3	1	$\frac{1}{3}$
$\log_{1/3}(x)$	-2	-1	0	1

(b)

**5.2.7.5.****Answer.**

(a) $x = 10,000$

(b) $x = 10^8$

5.2.7.7.**Answer.** $0 < x < 0.01$ **5.2.7.9.****Answer.**

(a)	$\log(100, 322) \approx 5.001$	(b)	$\log(693) \approx 2.841$
-----	--------------------------------	-----	---------------------------

5.2.7.11.**Answer.**

(a)	$\log(-7)$ is undefined.	(b)	$6 \log(28) \approx 8.683$
-----	--------------------------	-----	----------------------------

5.2.7.13.**Answer.**

(a) 15.614 (b) 0.419

5.2.7.15.**Answer.**

(a)	81	(c)	Definition of logarithm base 3	(d)	1.8
(b)	4	(e)	a		

5.2.7.17.**Answer.**

(a) 2^8 (b) -2

5.2.7.19.**Answer.**

(a)	$2k$	(c)	\sqrt{x}
(b)	x^3	(d)	$2m$

5.2.7.21.**Answer.**

(a) $(9, \infty)$ (b) $f^{-1}(x) = 3^{x-4} + 9$

5.2.7.23.**Answer.**

(a) $f^{-1}(x) = \log_4(100 - x) - 2$

(b) $f^{-1}(f(1)) = f^{-1}(36) = \log_4(64) - 2 = 1$

(c) $f(f^{-1}(84)) = f(0) = 100 - 4^2 = 84$


5.2.7.25.**Answer.**

(a) IV

(b) I

(c) II

(d) III

5.2.7.27.**Answer.**

(b) The graph resembles a logarithmic function. The (translated) log function is close to the points but appears too steep at first and not steep enough after $n = 15$. Overall, it is a good fit.

(c) f grows (more and more slowly) without bound. f will eventually exceed 100 per cent, but no one can forget more than 100% of what is learned.

5.2.7.29.**Answer.**

(a) $10^{1.41} \approx 25.704$

(c) $10^{0.52} \approx 3.3113$

(b) $10^{-1.69} \approx 0.020417$

5.2.7.31.**Answer.** $16^w = 256$ **5.2.7.33.****Answer.** $b^{-2} = 9$ **5.2.7.35.****Answer.**
 $10^{-2.3} = A$ **5.2.7.37.****Answer.** $u^w = v$ **5.2.7.39.****Answer.** $b = 2$ **5.2.7.41.****Answer.** $b = 100$ **5.2.7.43.****Answer.** $x = 11$ **5.2.7.45.****Answer.** $x = 7^{2/3}$ **5.2.7.47.****Answer.** $x = 4$ **5.2.7.49.****Answer.** $x = 11$ **5.2.7.51.****Answer.** $x = 3$ **5.2.7.53.****Answer.** No solution

5.2.7.55.**Answer.** $A = k(10^{t/T} - 1)$ **5.2.7.57.****Answer.** $s = \frac{b^{N/N_0}}{k}$ **5.2.7.59.****Answer.** $H = (H_0)^{kM^2}$ **5.2.7.61.****Answer.**

(a) II


(c) III

(e) I


(b) VI

(d) V


(f) IV

5.2.7.63.**Answer.**


No inverse function



No inverse function

5.2.7.65.**Answer.**

The functions are equal.

5.2.7.67.**Answer.**

The functions are equal.

5.2.7.69.**Answer.**

(a)


x	x^2	$\log(x)$	$\log(x^2)$
1	1	0	0
2	4	0.301	0.602
3	9	0.477	0.954
4	16	0.602	1.204
5	25	0.699	1.398
6	36	0.778	1.556

(b) $\log(x^2) = 2 \log(x)$

5.2.7.71.

Answer.

x	$y = \log_e(x)$
1	0
2	0.693
4	1.386
16	2.772
$\frac{1}{2}$	-0.693
$\frac{1}{4}$	-1.386
$\frac{1}{16}$	-2.772



5.3 · The Natural Base

5.3.8 · The Natural Logarithm (Homework 5.3)



5.3.8.1.

Answer.

5.3.8.3.

Answer.

x	-10	-5	0	5	10	x	15	10	20	-10	20	-5	0	5	10	15	20
$f(x)$	0.135	0.368	1	2.718	7.389	e^x	20.086	54.598	148.413	0.000045	0.0000002	0.0000000007	1	0.223	0.05	0.011	0.00248



5.3.8.5.

Answer.

(a) 2

(b) $5t$

(c) $\frac{1}{x}$

(d) $\frac{1}{2}$

5.3.8.7.

Answer.

(a) 0.64

(b) 3.81

(c) -1.20

5.3.8.9.

Answer.

(a) 4.14

(b) 1.88


(c) 0.07

5.3.8.11.**Answer.**

(a) $N(t) = 6000e^{0.04t}$

(b)	t	0	5	10	15	20	25	30
	$N(t)$	6000	7328	8951	10,933	13,353	16,310	19,921

(c)




(d) 15,670

(e) 70.3 hrs

5.3.8.13.**Answer.**

(a)



(b) 941.8 lumens

(c) 2.2 cm

5.3.8.15.**Answer.**

$$P(t) = 20(e^{0.4})^t \approx 20 \cdot 1.492^t;$$

increasing; initial value 20

5.3.8.17.**Answer.**

$$P(t) = 6500(e^{-2.5})^t \approx 6500 \cdot 0.082^t;$$

decreasing; initial value 6500

5.3.8.19.**Answer.**

(a)	x	0	0.5	1	1.5	2	2.5
	e^x	1	1.6487	2.7183	4.4817	7.3891	12.1825

(b) Each ratio is $e^{0.5} \approx 1.6487$: Increasing x -values by a constant $\Delta x = 0.5$ corresponds to multiplying the y -values of the exponential function by a constant factor of $e^{\Delta x}$.

5.3.8.21.**Answer.**

(a)	x	0	0.6931	1.3863	2.0794	2.7726	3.4657	4.1589
	e^x	1	2	4	8	16	32	64

(b) Each difference in x -values is approximately $\ln(2) \approx 0.6931$: Increasing x -values by a constant $\Delta x = \ln(2)$ corresponds to multiplying the y -values of the exponential function by a constant factor of $e^{\Delta x} = e^{\ln(2)} = 2$. That

is, each function value is approximately equal to double the previous one.

5.3.8.23.

Answer. 0.8277

5.3.8.27.

Answer. 1.6451

5.3.8.31.

Answer. $t = \frac{1}{k} \ln(y)$

5.3.8.35.

Answer. $k = e^{T/T_0} - 10$

5.3.8.25.

Answer. -2.9720

5.3.8.29.

Answer. -3.0713

5.3.8.33.

Answer. $t = \ln\left(\frac{k}{k-y}\right)$

5.3.8.37.

Answer.

(a)

n	0.39	3.9	39	390
$\ln(n)$	-0.942	1.361	3.664	5.966

(b) Each difference in function values is approximately $\ln(10) \approx 2.303$: Multiplying x -values by a constant factor of 10 corresponds to adding a constant value of $\ln 10$ to the y -values of the natural log function.

5.3.8.39.

Answer.

(a)

n	2	4	8	16
$\ln(n)$	0.693	1.386	2.079	2.773


(b) Each quotient equals k , where $n = 2^k$. Because $\ln(n) = \ln(2^k) = k \cdot \ln(2)$,

$$k = \frac{\ln(n)}{\ln(2)}.$$

5.3.8.41.

Answer.


(a)
$$N(t) = 100e^{(\ln(2))t} \approx 100e^{0.6931t}$$



5.3.8.43.

Answer.


(a)
$$N(t) = 1200e^{(\ln(0.6))t} \approx 1200e^{-0.5108t}$$



5.3.8.45.

Answer.

(a)
$$N(t) = 10e^{(\ln(1.15))t} \approx 10e^{0.1398t}$$



5.3.8.47.

Answer.

(a) 20,000

(b) $\left(\frac{35,000}{20,000}\right)^{1/10} \approx e^{0.056}$

(c) $P(t) = 20,000e^{0.056t}$

(d) 107,188

5.3.8.49.

Answer.

(a) $\left(\frac{385}{500}\right)^{1/2} \approx e^{-0.1307}$

(b) $N(t) = 500e^{-0.1307t}$


(c) 135.3 mg

5.3.8.51.

Answer.

- (a) $A(t) = 500e^{0.095t}$ (b) 7.3 years (c) 7.3 years


d-e



5.3.8.53.

Answer.

- (a) 6 hours




- (b) 6 hours

5.3.8.55.

Answer.


- (a) $\frac{1}{2}N_0, \frac{1}{4}N_0, \frac{1}{16}N_0$



- (c) $N(t) = N_0e^{-0.0866t}$

5.3.8.57.

Answer.



- (a)




$$y = 116(0.975)^t$$

- (b) $G(t) = 116e^{-0.025t}$

- (c) 28 minutes

5.4 · Logarithmic Scales

5.4.8 · Logarithmic Scales (Homework 5.4)

5.4.8.1.**Answer.****5.4.8.3.****Answer.****5.4.8.5.****Answer.** 1.58, 6.31, 15.8, 63.1**5.4.8.7.****Answer.** 1, 80, 330, 1600, 7000, 4×10^7 **5.4.8.9.****Answer.****5.4.8.11.****Answer.** Proxima Centauri: 15.5; Barnard: 13.2; Sirius: 1.4; Vega: 0.6; Arcturus: -0.4; Antares: -4.7; Betelgeuse: -7.2**5.4.8.13.****Answer.**

- | | |
|------------|--------------------------|
| (a) 1 | (e) 0.000079 |
| (b) 0.5012 | (f) 3.2×10^{-7} |
| (c) 0.1259 | (g) 2×10^{-8} |
| (d) 0.01 | (h) 8×10^{-10} |

5.4.8.15.**Answer.**

- | | |
|---------------------------------|--|
| (a) $10^{1.75} \approx 56.2341$ | (b) $10^{(\log(600))/2} \approx 24.4949$ |
|---------------------------------|--|

5.4.8.17.**Answer.** $10^{3.4} \approx 2512$ **5.4.8.19.****Answer.** A: $a \approx 45$, $p \approx 7.4\%$; B: $a \approx 400$, $p \approx 15\%$; C: $a \approx 6000$, $p \approx 50\%$; D: $a \approx 13000$, $p \approx 45\%$ **5.4.8.21.****Answer.** 3.2**5.4.8.23.****Answer.** 0.0126

5.4.8.25.

Answer. 100

5.4.8.27.

Answer. 6,309,573 watts per square meter

5.4.8.29.

Answer. 1000

5.4.8.31.

Answer. 12.6

5.4.8.33.

Answer. 100

5.4.8.35.Answer. $\approx 25,000$ **5.4.8.37.**

Answer. 4.7

5.4.8.39.

Answer. 53

5.5 · Chapter Summary and Review**5.5.2 · Chapter 5 Review Problems****5.5.2.1.**

Answer.

y	-1	1	3	11
$x = f^{-1}(y)$	-1	0	1	2

5.5.2.3.

Answer.

y	0	$-\frac{1}{3}$	-1	-3
$w = g^{-1}(y)$	-1	0	1	2


5.5.2.5.

Answer.

$$(a) P^{-1}(350) = 40 \qquad (b) P^{-1}(100) = 0$$


5.5.2.7.

Answer.

$$(a) f^{-1}(x) = x - 4 \qquad (b) $$


5.5.2.9.

Answer.

$$(a) f^{-1}(x) = \sqrt[3]{x+1} \qquad (b) $$

5.5.2.11.

Answer.

$$(a) f^{-1}(x) = \frac{1}{x-2} \qquad (b) $$

5.5.2.13.

Answer. 0

5.5.2.15.

Answer.

(a) $f^{-1}(300) = 200$: \$200,000 in advertising results in \$300,000 in revenue.

(b) $f(A) = 250$ or $A = f^{-1}(250)$

5.5.2.17.

Answer. $10^z = 0.001$

5.5.2.21.

Answer. $b^3 = 3x + 1$

5.5.2.25.

Answer. $6n$

5.5.2.29.

Answer. -1

5.5.2.35.

Answer. $\frac{-15}{8}$

5.5.2.37.

Answer. $\frac{9}{4}$

5.5.2.39.

Answer. 3

5.5.2.41.

Answer. $x \approx 1.548$

5.5.2.43.

Answer. $x \approx 411.58$

5.5.2.45.

Answer. $x \approx 2.286$

5.5.2.47.

Answer. \sqrt{x}

5.5.2.49.

Answer. $k - 3$

5.5.2.51.

Answer.

(a) $P = 7,894,862e^{-0.011t}$

(b) 1.095%

5.5.2.53.

Answer.

(a) \$1419.07

(b) 13.9 years

(c) $t = 20 \ln \left(\frac{A}{1000} \right)$

5.5.2.55.

Answer. $t = \frac{-1}{k} \ln \left(\frac{y-6}{12} \right)$

5.5.2.57.

Answer. $M = N^{Qt}$

5.5.2.59.


Answer. $P(t) = 750(1.3771)^t$

5.5.2.61.

Answer. $N(t) = 600e^{-0.9163t}$

5.5.2.63.

Answer.



5.5.2.65.

Answer. Order 3: 17,000; Order 4: 5000; Order 8: 40; Order 9: 11

5.5.2.67.

Answer. 5×10^{-7}

5.5.2.69.

Answer. 3160

6 · Quadratic Functions


6.1 · Factors and x -Intercepts

6.1.7 · Factors and x -intercepts (Homework 6.1)

6.1.7.1.

Answer.

(a)	t	0	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
	h	300	306	304	294	276	250	216	174	124	66	0



(c) 306.25 ft at 0.625 sec

(d) 1.25 sec

(e) 5 sec

6.1.7.3.

Answer. $-\frac{5}{2}, 2$

6.1.7.7.

Answer. $-\frac{3}{4}, -8$

6.1.7.11.

Answer. $\frac{1}{2}, -3$

6.1.7.15.

Answer. 1

6.1.7.19.

Answer. 2, 3

6.1.7.23.

Answer. $-3, 6$

6.1.7.5.

Answer. $0, \frac{-10}{3}$

6.1.7.9.

Answer. 4

6.1.7.13.

Answer. 0, 3

6.1.7.17.

Answer. $\frac{1}{2}, 1$

6.1.7.21.

Answer. $-1, 2$

6.1.7.25.

Answer. The 3 graphs have the same x -intercepts. In general, the graph of $y = ax^2 + bx + c$ has the same x -intercepts as the graph of $y = k(ax^2 + bx + c)$.

6.1.7.27.

Answer. The 3 graphs have the same x -intercepts. In general, the graph of $y = ax^2 + bx + c$ has the same x -intercepts as the graph of $y = k(ax^2 + bx + c)$.

6.1.7.29.**Answer.**

$$x^2 + x - 2 = 0$$

6.1.7.31.**Answer.**

$$x^2 + 5x = 0$$

6.1.7.33.**Answer.**

$$2x^2 + 5x - 3 = 0$$

6.1.7.35.**Answer.**

$$8x^2 - 10x - 3 = 0$$

6.1.7.37.**Answer.**

$$f(x) = 0.1(x - 18)(x + 15)$$

6.1.7.39.**Answer.**

$$g(x) = -0.08(x - 18)(x + 32)$$

6.1.7.41.**Answer.**

(a) $10^2 + h^2 = (h + 2)^2$

(b) 24 ft

6.1.7.43.**Answer.**


(a) $h = -16t^2 + 16t + 8$

(b) 12 ft; 8 ft

(c) $11 = -16t^2 + 16t + 8$; at $\frac{1}{4}$ sec and $\frac{3}{4}$ sec

(d) $\Delta T_{bl} = 0.25$

(e)



(f) 1.37 sec

6.1.7.45.**Answer.**

Width	Length	Area
10	170	1700
20	160	3200
30	150	4500
40	140	5600
50	130	6500
60	120	7200
70	110	7700
80	100	8000

(a)

(b) $l = 180 - x$, $A = 180x - x^2$; 80 yd by 100 yd(c) $180x - x^2 = 8000$, 80 yd by 100 yd, or 100 yd by 80 yd. There are two solutions because the pasture can be oriented in two directions.**6.1.7.47.****Answer.**(a) $l = x - 4$, $w = x - 4$, $h = 2$, $V = 2(x - 4)^2$

(b)

x	4	5	6	7	8	9	10
V	0	2	8	18	32	50	72

(c) As x increases, V increases.

(d) 9 inches by 9 inches.

(e) $2(x - 4)^2 = 50$, $x = 9$ **6.1.7.49.****Answer.**


(a)

x	0	500	1000	1500	2000	2500	3000	3500
I	0	550	1000	1350	1600	1750	1800	1750

x	4000	4500	5000	5500	6000	6500	7000
I	1600	1350	1000	550	0	-650	-1400

(b) 1600, 1000, -1400

(c)



(d) No increase

(e) 3000; 1800

6.1.7.51.**Answer.** ± 1 **6.1.7.53.****Answer.** $\sqrt[3]{-3/4}$, 1

6.1.7.55.Answer. $-27, 1$ **6.1.7.59.**Answer. $1, 2$ **6.1.7.57.**Answer. $\log(2), \log(3)$ **6.1.7.61.**Answer. $\frac{-1}{6}, 1$ **6.1.7.63.**

Answer.

(a) $A = \frac{1}{2}(x^2 - y^2)$

(b) $A = \frac{1}{2}(x - y)(x + y)$

(c) 18 sq ft

6.2 · Solving Quadratic Equations**6.2.8 · Solving Quadratic Equations (Homework 6.2)****6.2.8.1.**

Answer.

(a) $(x + 4)^2$ (b) $\left(x - \frac{7}{2}\right)^2$ (c) $\left(x + \frac{3}{4}\right)^2$ (d) $\left(x - \frac{2}{5}\right)^2$

6.2.8.3.Answer. 1 **6.2.8.7.**

Answer. $\frac{3}{2} \pm \sqrt{\frac{21}{4}} = \frac{-3 \pm \sqrt{21}}{2}$

6.2.8.11.

Answer. $\frac{-4}{3}, 1$

6.2.8.15.

Answer. $-1, \frac{4}{3}$

6.2.8.19.

Answer. $-1 \pm \sqrt{1 - c}$

6.2.8.21.

Answer.
$$\frac{-\frac{b}{2} \pm \sqrt{\frac{b^2 - 4}{4}}}{2} = \frac{-b \pm \sqrt{b^2 - 4}}{4}$$

6.2.8.23.

Answer.
$$\frac{-1 \pm \sqrt{4a + 1}}{a}$$


6.2.8.25.

Answer.

(a) $A = (x + y)^2$

(b) $A = x^2 + 2xy + y^2$

(c) x^2, xy, xy, y^2

**6.2.8.27.**Answer. $1.618, -0.618$ **6.2.8.29.**Answer. $1.449, -3.449$

6.2.8.31.

Answer. 1.695, -0.295

6.2.8.35.

Answer. -5.894, 39.740

6.2.8.33.


Answer. 1.434, 0.232

6.2.8.37.

Answer.

(a)

s	10	20	30	40	50	60	70	80	90	100
d	9	27	53	87	129	180	239	307	383	467




(c) $\frac{s^2}{24} + \frac{s}{2} = 50$; 29.16 mph

6.2.8.39.

Answer.

(a)

t	0	5	10	15	20	25
h	11,000	10,520	9240	7160	4280	600



(c) $-16t^2 - 16t + 11,000 = 1000$; 24.5 sec

(d) 1.2 sec

6.2.8.41.

Answer.

(a) $2l + 4w = 100$

(b) $l = 50 - 2w$

(c) $w(50 - 2w) = 250$; $w = 6.91, 18.09$

(d) 12.06 m by 6.91 m, or 4.61 m by 18.09 m

6.2.8.43.

Answer.

(a) 47.2 mi

(b) 1.26 mi

6.2.8.45.

Answer.

$$w = \frac{-4l \pm \sqrt{16l^2 + 8A}}{4} = \frac{-2l \pm \sqrt{4l^2 + 2A}}{2}$$

6.2.8.47.

Answer.

$$t = \frac{4 \pm \sqrt{16 + 64h}}{32} = \frac{1 \pm \sqrt{1 + 4h}}{8}$$

6.2.8.49.

$$\text{Answer. } t = \frac{v \pm \sqrt{v^2 - 2as}}{a}$$

6.2.8.51.

$$\text{Answer. } y = \frac{-x \pm \sqrt{8 - 11x^2}}{2}$$

6.2.8.53.

$$\text{Answer. } 0, x^2$$

6.2.8.55.

$$\text{Answer. } \frac{-3x \pm 3}{2}$$

6.2.8.57.

$$\text{Answer. } \frac{\pm \sqrt{4x^2 - 36}}{3} = \frac{\pm 2\sqrt{x^2 - 9}}{3}$$

6.2.8.59.

$$\text{Answer. } \frac{\pm 2x}{5}$$

6.2.8.61.

$$\text{Answer. } 3 \pm \sqrt{\frac{V}{\pi h}}$$

6.2.8.63.

$$\text{Answer. } \pm \sqrt{\frac{2(E - mgh)}{m}}$$

6.2.8.65.

$$\text{Answer. } \pm \sqrt{\frac{V}{2w} - s^2}$$

6.2.8.67.

$$\text{Answer. } \frac{-b}{2a}$$

6.2.8.69.

$$\text{Answer. } \frac{-b \pm \sqrt{b^2 - 4c}}{2}$$


6.3 · Graphing Parabolas

6.3.9 · Graphing Parabolas (Homework 6.3)


6.3.9.1.

Answer.


- (a) The parabola opens up, twice as steep as the standard parabola.




- (b) The parabola is the standard parabola shifted 2 units up.



- (c) The parabola is the standard parabola shifted 2 units left.




(d) The parabola is the standard parabola shifted 2 units down.




6.3.9.3.

Answer.


a Vertex $(0, -16)$; x -intercepts $(\pm 4, 0)$




b Vertex $(0, 16)$; x -intercepts $(\pm 4, 0)$



c Vertex $(8, 64)$; x -intercepts $(0, 0)$ and $(16, 0)$




d Vertex $(8, -64)$; x -intercepts $(0, 0)$ and $(16, 0)$




6.3.9.5.

Answer.


a Vertex $(1, -3)$; x -intercepts $(0, 0)$ and $(-2, 0)$




b Vertex $(1, -3)$; x -intercepts $(0, 0)$ and $(2, 0)$



c Vertex $(0, 6)$; no x -intercepts



d Vertex $(0, -6)$; x -intercepts $(\pm\sqrt{2}, 0)$



6.3.9.7.


Answer.

(a) II (b) IV (c) I (d) III (e) VI (f) V

6.3.9.9.

Answer.

(a) $(2000, 400)$; The largest annual increase in biomass, 400 tons, occurs when the biomass is 2000 tons.



(c) $4000 < x \leq 5000$; When there are too many fish, there will not be enough food to support all of them.


6.3.9.11.

Answer.

(a)

x	1	2	3	4	5
y	1.6	1.2	0.8	0.4	0
A	1.6	2.4	2.4	1.6	0

(b) $A = x(2 - 0.4x)$ or $A = 2x - 0.4x^2$




(d) The maximum number of young marmots, on average, is 2.5; the optimal number of female marmots is 2.5.

6.3.9.13.

Answer.

(a)



Vertex: $(-6, -1.5)$; Horizontal intercepts $(-12, 0)$ and $(0, 0)$. The point $(0, 0)$ means that no distance is required to stop a plane that is not moving.

(b) 594 ft/sec

6.3.9.15.

Answer.

(a) $\left(\frac{3}{2}, \frac{17}{4}\right)$, maximum


(b) $\left(\frac{2}{3}, \frac{1}{9}\right)$, minimum

(c) $(-4.5, 18.5)$, maximum

6.3.9.17.

Answer.


(a) x -intercepts: $\left(\frac{-1}{2}, 0\right)$ and $(4, 0)$; y -intercept: $(0, 4)$; vertex: $\left(\frac{7}{4}, \frac{81}{8}\right)$



6.3.9.21.

Answer.


(a) No x -intercepts; y -intercept: $(0, 7)$; vertex: $(-2, 3)$



6.3.9.19.

Answer.


(a) x -intercepts: $(-2, 0)$ and $(1, 0)$; y -intercept: $(0, -1.2)$; vertex: $(-0.5, -1.35)$



6.3.9.23.



Answer.

(a) x -intercepts: $(-1 \pm \sqrt{2}, 0)$; y -intercept: $(0, -1)$; vertex: $(-1, -2)$



6.3.9.25.**Answer.**

(a) x -intercepts: $\left(\frac{3 \pm \sqrt{3}}{2}, 0\right)$;
 y -intercept: $(0, -3)$; vertex:
 $\left(\frac{3}{2}, \frac{3}{2}\right)$

**6.3.9.27.****Answer.**

$f(x) = x^2 - 6x + 5$: x -intercepts $(1, 0)$ and $(5, 0)$; $g(x) = x^2 - 6x + 9$:
 x -intercept $(3, 0)$; $h(x) = x^2 - 6x + 12$: No x -intercept.

- (b) $16, 0, -12$: $D = 16$ means that there are two rational x -intercepts, $D = 0$ means that there is exactly one x -intercept, $D = -12$ means that there is no x -intercept.

6.3.9.29.**Answer.** Two complex solutions**6.3.9.31.****Answer.** One repeated rational solution**6.3.9.33.****Answer.** Two distinct real solutions**6.3.9.35.****Answer.** No**6.3.9.37.****Answer.** Yes**6.3.9.39.****Answer.**

(a) $2 - \sqrt{5}$

6.3.9.41.**Answer.**

(a) $4 + 3\sqrt{2}$

(b) $x^2 - 4x - 1 = 0$

(b) $x^2 - 8x - 2 = 0$

6.3.9.43.**Answer.**

(a) IV

(b) V


(c) I

(d) VII

6.3.9.45.**Answer.**

(a) $y = x^2 + x - 6; x = \frac{-1}{2}$

(b) $y = 2x^2 + 2x - 12; x = \frac{-1}{2}$

6.3.9.47.**Answer.**


$(-1, -1), (-2, -4), (-3, -9), (-4, -16)$

(c) $y = -x^2$

(d) The vertex of $y = x^2 + 2kx$ is $(-k, -k^2)$

6.3.9.49.**Answer.**

(a)	t	0	0.5	1.0	1.5	2.0	2.5	3.0	3.5
	x	0	6.075	11.5	16.275	20.4	23.875	26.7	28.875
	y	0	7.44	12.48	15.12	15.36	13.2	8.64	1.68



(c) $y \approx 15.4$ m

(d) $x \approx 30$ m

(e) 3.6 sec

(f) $x \approx 29.2$ m

(g) $y \approx 15.55$ m

6.4 · Problem Solving**6.4.6 · Problem Solving (Homework 6.4)**

6.4.6.1.**Answer.**


No. of price increases	Price of room	No. of rooms rented	Total revenue
0	20	60	1200
1	22	57	1254
2	24	54	1296
3	26	51	1326
4	28	48	1344
5	30	45	1350
6	32	42	1344
7	34	39	1326
8	36	36	1296
10	40	30	1200
12	44	24	1056
16	52	12	624
20	60	0	0

(a)

(b) Price of a room: $20+2x$; Rooms rented: $60-3x$; Revenue: $1200+60x-6x^2$

(c) 20

(d)



(e) \$24; \$36

(f) \$1350; \$30; 45 rooms

6.4.6.3.**Answer.**

(a) (For example) 10 m by 20 m with area 200 sq m; or 15 m by 15 m, area 225 sq m

(b) $30 - x$ (c) $30x - x^2$ **6.4.6.5.****Answer.** 3 sec, 144 ft**6.4.6.7.****Answer.** 100 baskets, \$2000**6.4.6.9.****Answer.**(a) Length: $50 - w$; Area: $50w - w^2$

(b) 625 sq in

6.4.6.11.**Answer.**

(a) $300w - 2w^2$

(b) 11,250 sq yd

6.4.6.13.

Answer.

(a) Number of people: $16 + x$; Price per person: $2400 - 100x$; Total revenue: $38,400 + 800x - 100x^2$


(b) 20

6.4.6.15.

Answer. $a = 0.9$; $I = \$865.80$


6.4.6.17.

Answer.



6.4.6.19.

Answer.



6.4.6.21.


Answer.

(a) (3, 4)


6.4.6.23.

Answer.

(a) (-4, -3)



(c) $y = 2x^2 - 12x + 22$



(c) $y = -\frac{1}{2}x^2 - 4x - 11$

6.4.6.25.

Answer.

(a) $y = (x - 2)^2 + 3$

6.4.6.27.


Answer.

(a) $y = 3(x + 1)^2 - 5$


6.4.6.29.

Answer.


(a) $y = -2(x + 2)^2 + 11$



(b)




(b)




(b)

6.4.6.31.


Answer. No solutions:



One solution:



Two solutions:



6.4.6.33.

Answer. $(-1, 12), (4, 7)$

6.4.6.37.

Answer. No solution

6.4.6.41.

Answer. $(1, 4)$

6.4.6.45.

Answer.

6.4.6.35.

Answer. $(-2, 7)$


6.4.6.39.

Answer. $(-2, -5), (5, 16)$

6.4.6.43.

Answer. $(3, 1)$

(a)



(b) Larger, by 75 tons. Smaller, by 125 tons.


(c) 1000 tons and 3000 tons

(d) The fish population will decrease each year until it is completely depleted.

6.4.6.47.

Answer.

(a)



(b) $K > N$. The population will decrease by 48 bears.

(c) The population will increase by 18 bears.

(d) 1000

(e) Populations between 0 and 1000 will increase; populations over 1000 will decrease.


(f) 1000 (unless the population is 0)

(g) 500 (unless the population is 0)

6.4.6.49.

Answer.


(a) (200, 2600), (1400, 18,200) (c) $x = 800$



6.4.6.51.

Answer.


(a) (60, 39,000), (340, 221,000)



(c) $x = 200$

6.4.6.53.

Answer.



(b) See graph and (c)

(c) $ad^2 + k$

(d) The two points on the parabola that are the same horizontal distance from the line $x = h$ the axis of symmetry have the same y -coordinate, so they are symmetric about that line.

6.5 · Chapter Summary and Review

6.5.2 · Chapter 6 Review Problems

6.5.2.1.

Answer. $0, \frac{-5}{2}$

6.5.2.3.

Answer. $-1, 2$

6.5.2.5.

Answer. $-2, 3$

6.5.2.7.


Answer. $4x^2 - 29x - 24 = 0$

6.5.2.9.**Answer.** $y = (x - 3)(x + 2.4)$ **6.5.2.11.****Answer.** 1, 2**6.5.2.13.****Answer.** $-1, \frac{1}{4}$ **6.5.2.15.****Answer.** $2 \pm \sqrt{10}$ **6.5.2.17.****Answer.** $\frac{3 \pm \sqrt{3}}{2}$ **6.5.2.19.****Answer.** 1, 2**6.5.2.21.****Answer.** $2 \pm \sqrt{2}$ **6.5.2.23.****Answer.** $\pm \sqrt{\frac{2K}{m}}$ **6.5.2.25.****Answer.** $\frac{3 \pm \sqrt{9 - 3h}}{3}$ **6.5.2.27.****Answer.** 9**6.5.2.29.****Answer.** 10 ft by 18 ft or 12 ft by 15 ft**6.5.2.31.****Answer.** 1 sec**6.5.2.33.****Answer.**

(a) $h = 100t - 2.8t^2$

(c) 893 ft


(b)

(d) $15\frac{5}{7}$ sec on the way up and 20 sec on the way down**6.5.2.35.****Answer.** A_1 is the area of a square minus the area of two triangles:


$$x^2 - 2 \left(\frac{1}{2} y \cdot y \right) = x^2 - y^2$$

6.5.2.37.**Answer.**


- (a) Vertex and intercepts are all $(0, 0)$.

**6.5.2.41.****Answer.**


- (a) Vertex $(\frac{-1}{2}, \frac{-25}{4})$; x -intercepts $(-3, 0)$ and $(2, 0)$; y -intercept $(0, -6)$

**6.5.2.45.****Answer.**


- (a) Vertex $(\frac{1}{2}, \frac{-37}{4})$; x -intercepts $(\frac{1 \pm \sqrt{37}}{2}, 0)$; y -intercept $(0, -9)$

**6.5.2.47.****Answer.** Two**6.5.2.49.****Answer.** One rational solution**6.5.2.39.****Answer.**


- (a) Vertex $(\frac{9}{2}, \frac{-81}{4})$; x -intercepts $(9, 0)$ and $(0, 0)$; y -intercept $(0, 0)$

**6.5.2.43.****Answer.**

- (a) Vertex $(\frac{-1}{4}, \frac{65}{8})$; x -intercepts $(\frac{-1 \pm \sqrt{65}}{4}, 0)$; y -intercept $(0, 8)$

**6.5.2.53.****Answer.****6.5.2.51.****Answer.** No real solutions

- (a) 45; \$410
- (b)



6.5.2.55.


Answer.

- (a) $y = 60(4 + x)(32 - 4x)$
- (b) 2

6.5.2.57.

Answer.


- (a) $0, \frac{-3}{2}$
- (b) $\frac{5}{6}$
- (c) $-5, \frac{1}{2}$



6.5.2.59.


Answer.

- (a) 0, 1
- (b) None
- (c) -1, 3




6.5.2.61.

Answer. $(-\infty, -2) \cup (3, \infty)$




6.5.2.63.

Answer. $\left[-1, \frac{3}{2}\right]$



6.5.2.65.

Answer. $[-2, 2]$




6.5.2.67.

Answer.

(a) $R = p \left(220 - \frac{1}{4}p \right)$


6.5.2.69.

Answer. (1, 3), (-1, 3)



6.5.2.73.

Answer. (-9, 155), (5, 15)



6.5.2.77.

Answer. $a = 1, b = -1, c = -6$

6.5.2.79.

Answer. $p(x) = \frac{-1}{2}x^2 - 4x + 10$


6.5.2.81.

Answer. $y = 0.2(x - 15)^2 - 6$

6.5.2.83.

Answer.

(a) $f(x) = (x - 12)^2 - 100$




6.5.2.87.

Answer.

(a) $h = 36.98t + 5.17$

(b) 116.1 m, 153.1 m




(d) $h = -4.858t^2 + 47.67t + 0.89$

(b) Between \$4.00 and \$4.80


6.5.2.71.

Answer. (-1, -4), (5, 20)



6.5.2.75.


Answer. (-1, 2), (3, 0)




6.5.2.85.

Answer.

(a) $y = \frac{1}{3}(x + 3)^2 - 2$



(e) 100.2 m, 113.9 m



(g) Quadratic: Gravity will slow the cannonball, giving the graph a concave down shape.

7 · Polynomial and Rational Functions

7.1 · Polynomial Functions

7.1.6 · Polynomial Functions (Homework 7.1)

7.1.6.1.

Answer. $12x^3 - 5x^2 - 8x + 4$

7.1.6.5.

Answer. $6a^4 - 5a^3 - 5a^2 + 5a - 1$

7.1.6.3.

Answer. $x^3 - 6x^2 + 11x - 6$

7.1.6.7.

Answer. $y^4 + 5y^3 - 20y - 16$

7.1.6.9.

Answer. $6 + x + 5x^2$

7.1.6.11.

Answer. $4 - 7x^2 - 8x^4$

7.1.6.13.

Answer. $0x^2$

7.1.6.15.

Answer. $-8x^3$

7.1.6.17.

Answer.

(a) 4

(b) 5

(c) 7

7.1.6.19.

Answer. $(x + y)^3 = (x + y)(x + y)^2$
 $= (x + y)(x^2 + 2xy + y^2)$
 $= x^3 + 2x^2y + xy^2 + x^2y + 2xy^2 + y^3$
 $= x^3 + 3x^2y + 3xy^2 + y^3$

7.1.6.21.

Answer. $(x + y)(x^2 - xy + y^2) = x^3 - x^2y + xy^2 + x^2y - xy^2 + y^3$
 $= x^3 + y^3$

7.1.6.23.

Answer.

- (a) The formula begins with x^3 and ends with y^3 . As you proceed from term to term, the exponents on x decrease while the exponents on y increase, and on each term the sum of the exponents is 3. The coefficients of the two middle terms are both 3.
- (b) The formula is the same as for $(x - y)^3$, except that the terms alternate in sign.

7.1.6.25.

Answer. $1 + 6z + 12z^2 + 8z^3$

7.1.6.27.

Answer. $1 - 15\sqrt{t} + 75t - 125t\sqrt{t}$

7.1.6.29.**Answer.** $x^3 - 1$ **7.1.6.33.****Answer.** $27a^3 - 8b^3$ **7.1.6.31.****Answer.** $8x^3 + 1$ **7.1.6.35.****Answer.**
 $(x + 3)(x^2 - 3x + 9)$ **7.1.6.41.****Answer.** $(3a + 4b)(9a^2 - 12ab + 16b^2)$ **7.1.6.37.****Answer.**
 $(a - 2b)(a^2 + 2ab + 4b^2)$ **7.1.6.43.****Answer.** $(5ab - 1)(25a^2b^2 + 5ab + 1)$ **7.1.6.39.****Answer.** $(xy^2 - 1)(x^2y^4 + xy^2 + 1)$ **7.1.6.45.****Answer.**
 $(4t^3 + w^2)(16t^6 - 4t^3w^2 + w^4)$ **7.1.6.47.****Answer.**

(a) $\left(6 - \frac{5}{4}\pi\right)x^2$

(b) ≈ 132.67 square inches

7.1.6.49.**Answer.**

(a) $\frac{2}{3}\pi r^3 + \pi r^2 h$

(b) $V(r) = \frac{14}{3}\pi r^3$

7.1.6.51.**Answer.**

(a) $500(1 + r)^2$; $500(1 + r)^3$; $500(1 + r)^4$

(b) $500r^2 + 1000r + 500$; $500r^3 + 1500r^2 + 1500r + 500$; $500r^4 + 2000r^3 + 3000r^2 + 2000r + 500$

(c) \$583.20, \$629.86, \$680.24

7.1.6.53.**Answer.**

(a) Length: $16 - 2x$; Width: $12 - 2x$; Height: x


(b) $V = x(16 - 2x)(12 - 2x)$

(c) Real numbers between 0 and 6

(d)

x	1	2	3	4	5
V	140	192	180	128	60

(e)




(f) 2.26 in, 194.07 cu in

7.1.6.55.**Answer.**

(a) 0, 9

(b) $0 \leq x \leq 9$; $R \geq 0$ for these values

(c)


(d) $\frac{28}{3}$ points

(e) 36 points

(f) 3 ml or 8.2 ml

7.1.6.57.**Answer.**

(a)




(b) 900; 11, 145; 15, 078

(c) 1341; 171; 627

(d) Between 1990 and 1991

7.1.6.59.**Answer.**

(a)

(b) The graph is concave down until about $x = 12.5$ and is concave up afterwards. The cost is growing at the slowest rate at the inflection point at about $x = 12.5$, or 1250 students.


(c) About 2890

7.1.6.61.**Answer.**

(a) 20 cm

(b) 100 cm

7.1.6.63.**Answer.**(a) $763.10 < H(t) < 864$




- (c) 864 min
 (d) 859.8 min
 (e) Within 34 days of the summer solstice
 (f) More than 66 days from the summer solstice

7.2 · Graphing Polynomial Functions

7.2.7 · Graphing Polynomial Functions (Homework 7.2)

7.2.7.1.


Answer.



- (a) The end behavior is the same as for the basic cubic because the lead coefficient is positive.
 (b) There is one x -intercept, no turning points, one inflection point.

7.2.7.5.


Answer.



- (a) The end behavior is the same as for the basic cubic because the lead coefficient is positive.
 (b) There are three x -intercepts, two turning points, one inflection point.

7.2.7.3.


Answer.






- (a) The end behavior is the opposite to the basic cubic (the graph starts in the upper left and extends to the lower right) because the lead coefficient is negative.
 (b) There is one x -intercept, no turning points, one inflection point.

7.2.7.7.


Answer.




- (a) The end behavior is the same as for the basic cubic because the lead coefficient is positive.
 (b) There are three x -intercepts, two turning points, one inflection point.

7.2.7.9.**Answer.**

(b) and (c) are the same.

7.2.7.11.**Answer.**


- (a) The end behavior is the same as for the basic quartic because the lead coefficient is positive.
- (b) There are two x -intercepts, one turning point, no inflection point.

7.2.7.15.**Answer.**


- (a) The end behavior is the same as for the basic quartic because the lead coefficient is positive.
- (b) There are two x -intercepts, one turning point, two inflection points.

7.2.7.19.

Answer. The graph of a cubic polynomial with a positive lead coefficient will have the same end behavior as the basic cubic, and a cubic with a negative lead coefficient will have the opposite end behavior. Each graph of a cubic polynomial has one, two, or three x -intercepts, it has two, one or no turning point, and it has exactly one inflection point.

7.2.7.13.**Answer.**

- (a) The end behavior is the opposite of the basic cubic (the graph starts in the lower left and ends in the lower right) because the lead coefficient is negative.
- (b) There are no x -intercepts, three turning points, two inflection points.


7.2.7.17.**Answer.**

- (a) The end behavior is the opposite of the basic cubic (the graph starts in the lower left and ends in the lower right) because the lead coefficient is negative.
- (b) There are no x -intercepts, one turning point, two inflection points.

7.2.7.21.

Answer.

(a)



$$(-2, 0), (-1, 0), (3, 0)$$


(b) $P(x) = (x + 2)(x + 1)(x - 3)$

(c) Yes

7.2.7.25.

Answer.

(a)



$$(-2, 0), (1, 0), (4, 0)$$


(b) $p(x) = (x + 2)(x - 1)(x - 4)$

(c) Yes

7.2.7.23.

Answer.

(a)



$$(-2, 0), (0, 0), (1, 0), (2, 0)$$


(b) $R(x) = (x + 2)(x)(x - 1)(x - 2)$

(c) Yes

7.2.7.27.

Answer.

(a)




$$(-2, 0), (2, 0), (3, 0)$$

(b) $r(x) = (x + 2)^2(x - 2)(x - 3)$

(c) Yes


7.2.7.29.

Answer.




7.2.7.33.

Answer.




7.2.7.31.

Answer.



7.2.7.35.


Answer.



7.2.7.37.**Answer.**


(a) 0 (multiplicity 2)

(b)


**7.2.7.39.****Answer.**

(a) 0 (multiplicity 2), 2 (multiplicity 2)


(b)

**7.2.7.41.****Answer.**(a) 0, ± 2


(b)

**7.2.7.43.****Answer.**(a) $\pm\sqrt{2}$, $\pm\sqrt{8}$

(b)

**7.2.7.45.****Answer.**(a) ± 1 , -3 (multiplicity 2)

(b)

**7.2.7.47.****Answer.**

$$P(x) = (x + 2)(x - 1)(x - 4)$$

7.2.7.51.


$$\text{Answer. } P(x) = (x - 2)^3(x + 2)$$

7.2.7.49.


$$\text{Answer. } P(x) = (x + 3)^2(x - 2)$$

7.2.7.53.**Answer.**


- (a) $y = x^3 - 4x + 3$; The graph of $y = f(x)$ shifted 3 units up.




- (b) $y = x^3 - 4x - 5$; The graph of $y = f(x)$ shifted 5 units down.




- (c) $y = (x - 2)^3 - 4(x - 2)$; The graph of $y = f(x)$ shifted 2 units right.




- (d) $y = (x + 3)^3 - 4(x + 3)$; The graph of $y = f(x)$ shifted 3 units left.

**7.2.7.55.****Answer.**


- (a) $y = x^4 - 4x^2 + 6$; The graph of $y = f(x)$ shifted 6 units up.




- (b) $y = x^4 - 4x^2 - 2$; The graph of $y = f(x)$ shifted 2 units down.



- (c) $y = (x - 1)^4 - 4(x - 1)^2$; The graph of $y = f(x)$ shifted 1 unit right.



- (d) $y = (x + 2)^4 - 4(x + 2)^2$; The graph of $y = f(x)$ shifted 2 units left.

**7.2.7.57.****Answer.** $q(x) = 2x^2 + 4x - 7$; $r(x) = -32$ **7.2.7.59.****Answer.** $q(x) = x^3 - 6x$; $r(x) = -6x + 5$ **7.2.7.61.****Answer.**

- (a) If $P(x)$ is a nonconstant polynomial with real coefficients and a is any real number, then there exist unique polynomials $q(x)$ and $r(x)$ such that

$$P(x) = (x - a)q(x) + r(x)$$

where $\deg r(x) < \deg(x - a)$.

- (b) Zero

- (c) $P(a) = (a - a)q(a) + r(a) = r(a)$. Because $\deg r(x) = 0$, $r(x)$ is a constant. That constant value is $P(a)$, so $P(x) = (x - a)q(x) + P(a)$.

7.2.7.63.**Answer.**

(a) From the remainder theorem, $P(x) = (x - a)Q(x) + P(a)$
 $= (x - a)Q(x) + 0$
 $= (x - a)Q(x)$

(b) By definition of a factor, if $x - a$ is a factor of $P(x)$, then $P(x) = (x - a)q(x)$, so $P(x) = (x - a)q(x) + 0$. The uniqueness guaranteed in the remainder theorem tells us that $P(a) = 0$.

7.2.7.65.

Answer.

(a) $P(1) = 0$

(b) $\frac{1 \pm \sqrt{5}}{2}$

7.2.7.67.

Answer.


(a) $P(-3) = 0$

(b) 0, 2, 4

7.2.7.69.

Answer.

(a) About $-1 < x < 2$




(b)

x	-1	-0.5	0	0.5	1	1.5	2
$f(x)$	0.368	0.607	1	1.649	2.718	4.482	7.389
$p(x)$	0.333	0.604	1	1.646	2.667	4.188	6.333

(c) 0.122

(d)



The error is relatively small for values of x between -3 and 2.5 .

7.3 · Complex Numbers

7.3.10 · Complex Numbers (Homework 7.3)

7.3.10.1.

Answer. $-4 + 5i$

7.3.10.3.

Answer. $-4 + i$

7.3.10.5.

Answer. $\frac{-5}{6} - \frac{\sqrt{2}}{6}i$

7.3.10.7.

Answer. $-3 \pm 2i$

7.3.10.9.

Answer. $\frac{1}{6} \pm \frac{\sqrt{11}}{6}i$

7.3.10.11.Answer. $13 + 4i$ **7.3.10.15.**Answer. $20 + 10i$ **7.3.10.19.**Answer. $46 + 14i\sqrt{3}$ **7.3.10.23.**Answer. $-2 - 2i$ **7.3.10.25.**Answer. $-1 + 4i$ **7.3.10.31.**Answer. $\frac{3}{4} - \frac{\sqrt{3}}{4}i$ **7.3.10.37.**

Answer.

(a) 0

(b) 0

7.3.10.41.

Answer.

(a) 0

(b) 0

7.3.10.43.Answer. $4z^2 + 49$ **7.3.10.47.**Answer. $v^2 - 8v + 17$ **7.3.10.13.**Answer. $-0.8 + 3.8i$ **7.3.10.17.**Answer. $-17 + 34i$ **7.3.10.21.**

Answer. 52

7.3.10.27.Answer. $7 + 4i$ **7.3.10.33.**Answer. $\frac{-2}{3} + \frac{\sqrt{5}}{3}i$ **7.3.10.29.**Answer. $\frac{-25}{29} + \frac{10}{29}i$ **7.3.10.35.**Answer. i **7.3.10.39.**

Answer.

(a) 0

(b) 0

7.3.10.45.Answer. $x^2 + 6x + 10$ **7.3.10.49.**Answer. $x \geq 5; x < 5$ **7.3.10.51.**

Answer.

(a) -1

(b) 1

(c) $-i$

(d) -1

7.3.10.53.

Answer.

(a) $2 - \sqrt{5}$ (b) $x^2 - 4x - 1$ **7.3.10.55.**

Answer.

(a) $4 + 3i$ (b) $x^2 - 8x + 25$ **7.3.10.57.**

Answer.

(a) 4


(b) 5

7.3.10.59.

Answer.
 $x^4 - 6x^3 + 23x^2 - 50x + 50$

7.3.10.63.


Answer.



The complex conjugates are reflections of each other across the real axis.

7.3.10.67.

Answer.



7.3.10.71.

Answer.


(a) $m = \frac{b}{a}$

7.3.10.61.

Answer.
 $x^4 - 7x^3 + 20x^2 - 19x + 13$

7.3.10.65.


Answer.



The complex conjugates are reflections of each other across the real axis.

7.3.10.69.

Answer.



(b) $m = \frac{a}{-b}$

(c) -1 ; The angle is 90° .

7.4 · Graphing Rational Functions

7.4.7 · Graphing Rational Functions (Homework 7.4)

7.4.7.1.

Answer.


(a) $t = \frac{150}{50 - v}$

(b)

v	0	5	10	15	20	25	30	35	40	45	50
t	3	3.33	3.75	4.29	5	6	7.5	10	15	30	—

The travel time increases as the headwind speed increases.

(c)




7.4.7.3.**Answer.**

(a) $0 \leq p < 100$

p	0	15	25	40	50	75	80	90	100
C	0	12.7	24	48	72	216	288	648	—

(c) 60%




(d) $p > 96\%$

(e) $p = 100$; As the percentage immunized approaches 100, the cost grows without bound.**7.4.7.5.****Answer.**

(a) $C = 8 + \frac{20,000}{n}$

n	100	200	400	500	1000	2000	4000	5000	8000
C	208	108	58	48	28	18	13	12	10.5

(c)



(d) 2000


(e) $n > 5000$

(f) $C = 8$; As n increases, the average cost per calculator approaches \$8.**7.4.7.7.****Answer.**


(a) $4500 + \frac{3000}{x}$; $C(x) = 6x + 4500 + \frac{3000}{x}$

x	10	20	30	40	50	60	70	80	90	100
C	4860	4770	4780	4815	4860	4910	5018	5073	5130	

(c) \$4768.33



(d) 22; 14



(e)

The graph of C approaches the line as an asymptote.

7.4.7.9.

Answer.


(a) The surface area is $2x^2 + 4xh = 96$. Solving for h , $h = \frac{96 - 2x^2}{4x} = \frac{24}{x} - \frac{x}{2}$.

(b) $V = 24x - \frac{1}{2}x^3$

(c)

x	1	2	3	4	5	6	7
h	23.5	11	6.5	4	2.3	1	-0.07
V	23.5	44	58.5	64	57.5	36	-3.5

If the base is more than 7 cm, the top and bottom alone exceed the total area allowed.




(d)

Maximum of 64 cu. cm

(e) 4 cm


(f) $h = 4$ cm



7.4.7.11.

Answer.

(a)	v	-100	-75	-50	-25	0	25	50	75	100
	P	338.15	358.92	382.41	409.19	440	475.83	518.01	568.4	629.66




(c) -20 m/sec; 68 m/sec

(d) $v > 12$ m/sec

(e) $v = 332$; As v approaches 332 m per sec, the pitch increases without bound.


7.4.7.13.

Answer.




7.4.7.17.

Answer.




7.4.7.21.

Answer.




7.4.7.15.

Answer.




7.4.7.19.

Answer.




7.4.7.23.

Answer.




7.4.7.25.

Answer.




7.4.7.29.

Answer.




7.4.7.27.

Answer.



7.4.7.31.

Answer.




7.4.7.33.

Answer.

$$a \quad y = \frac{2}{x} + 2$$

b




7.4.7.35.

Answer.

$$a \quad y = \frac{1}{x+1} + 1$$


b



7.4.7.37.

Answer.

$$\text{a } y = \frac{1}{(x-2)^2} + 3$$



7.4.7.39.

Answer.

$$\text{(a) } \frac{25}{s+8}$$

$$\text{(b) } \frac{25}{s-8}$$

$$\text{(c) } \frac{50s}{s^2-64}$$

7.4.7.41.

Answer.

$$\text{(a) } \frac{900}{400+w}$$

$$\text{(b) } \frac{900}{400-w}$$

$$\text{(c) Orville by } \frac{1800w}{160,000-w^2} \text{ hours}$$

7.4.7.43.

Answer.

$$\text{(a) } \frac{1}{f} = \frac{2q+60}{q^2+60q}$$


$$\text{(b) } f = \frac{q^2+60q}{2q+60}$$

7.4.7.45.

Answer.

$$\text{(a) } \frac{1}{y} = \frac{1}{x} + \frac{1}{k} = \frac{k+x}{xk}, \text{ so by taking reciprocals, } y = \frac{kx}{x+k}.$$

(b)



The graphs increase from the origin and approach a horizontal asymptote at $y = k$.


7.4.7.47.

$$\text{Answer. } \frac{12x}{x+20}$$

7.4.7.49.

Answer.

(a) V



$V \approx 0.7, K \approx 2.2$ (many answers are possible)

(b) $\frac{V}{2}$

(c)

(d) (See figure.)

7.4.7.51.


Answer.

(a) $\frac{1}{v} = \frac{K}{V} \cdot \frac{1}{s} + \frac{1}{V}$; Therefore, $a = \frac{K}{V}$ and $b = \frac{1}{V}$

(b)

$\frac{1}{s}$	3	1.5	1	0.6	0.4	0.3	0.15
$\frac{1}{v}$	12.5	7.1	5	3.3	2.6	2.2	1.7

(c)



$$\frac{1}{v} = 3.8 \cdot \frac{1}{s} + 1.1$$

(d) $V \approx 0.89, K \approx 3.37$


7.4.7.53.

Answer.

(a) $x \neq 2$

(b) $x + 2$

(c)




7.4.7.55.

Answer.

(a) $x \neq \pm 1$

(b) $\frac{1}{x-1}$

(c)



7.5 · Equations That Include Algebraic Fractions

7.5.5 · Equations that include Algebraic Fractions (Homework 7.5)

7.5.5.1.

Answer. $-\frac{1}{2}$

7.5.5.3.

Answer. $\frac{13}{8}$

7.5.5.5.

Answer. $\pm\sqrt{\frac{15}{8}}$

7.5.5.7.

Answer.
 $\frac{1800}{1849} \approx 0.97$


7.5.5.9.

Answer. 37 ft

7.5.5.11.

Answer.

(a) $t = \frac{150}{50 - v}$



(b) $4 = \frac{150}{50 - v}; v = 12.5$ mph


7.5.5.13.

Answer. $168 = \frac{72p}{100 - p}; p = 70\%$

7.5.5.15.

Answer.

(a)




(b) $x = 1$

7.5.5.17.

Answer.


(a)



(b) $x = \frac{1}{2}$

7.5.5.19.

Answer.



\$2.50

(b) $\frac{160}{x} = 6x + 49; x = 2.50$

7.5.5.21.

Answer.


(a) $L = \frac{3200}{w}$

Lowest point: (56.6, 226); The minimum perimeter is 226 ft for a width of 56.6 ft.

(b) $P = \frac{6400}{w} + 2w$

(d) $240 = \frac{6400}{w} + 2w$

(c)



(e) 40 ft by 80 ft

7.5.5.23.

Answer. Multiply both sides of the equation by bd and simplify.

$$\frac{a}{b} \cdot \frac{bc}{1} = \frac{c}{d} \cdot \frac{bc}{1}, \text{ so } ac = bd$$

7.5.5.25.

Answer. 4

7.5.5.27.

Answer. 40

7.5.5.29.

Answer. \$6187.50

7.5.5.31.

Answer. 45 mi

7.5.5.33.

Answer. 689

7.5.5.35.

Answer.

(a) 19,882 m

(b) 0.3%

(c) 0.00657 in

7.5.5.37.

Answer.

(a) $AE = 1$, $DE = x - 1$, $CD = 1$

(b) $\frac{1}{x} = \frac{x-1}{x}$

(c) $\frac{1 + \sqrt{5}}{2}$

7.5.5.39.

Answer. $r = \frac{S-a}{S}$

7.5.5.41.

Answer. $x = \frac{Hy}{2y-H}$

7.5.5.43.

Answer. $d = \pm \sqrt{\frac{Gm_1m_2}{F}}$

7.5.5.45.

Answer. $r = \frac{2QI}{I+Q}$

7.5.5.47.

Answer. $P = \frac{ES}{E+S}$

7.5.5.49.

Answer. 5

7.5.5.51.

Answer. 1

7.5.5.53.

Answer. $\frac{-14}{5}$

7.5.5.55.

Answer. $\frac{-1}{6}, \frac{-4}{3}$


7.5.5.57.

Answer.

(a) $t_1 = \frac{144}{s-20}$

(b) $t_2 = \frac{144}{s+20}$

(c)

**7.5.5.59.****Answer.**

(a) $t_1 = \frac{d}{r_1}, t_2 = \frac{d}{r_2}$

(c) $\frac{2d}{\frac{d}{r_1} + \frac{d}{r_2}}$

(b) Total distance is $2d$; total time $\frac{d}{r_1} + \frac{d}{r_2}$.

(d) $\frac{2r_1r_2}{r_1 + r_2}$

(e) $58\frac{1}{3}$ mph

7.6 · Chapter Summary and Review

7.6.2 · Chapter 7 Review Problems

7.6.2.1.

Answer. $2x^3 - 11x^2 + 19x - 10$

7.6.2.3.

Answer. $t^3 + 3t^2 - 5t - 4$

7.6.2.5.

Answer. $31x^2$

7.6.2.7.

Answer. $-13x^3$

7.6.2.9.

Answer. $(2x - 3z)(4x^2 + 6xz + 9z^2)$

7.6.2.11.

Answer. $(y + 3x)(y^2 - 3xy + 9x^2)$

7.6.2.13.

Answer. $v^3 - 30v^2 + 300v - 1000$

7.6.2.15.**Answer.**


(a) $\frac{1}{6}n^3 - \frac{1}{2}n^2 + \frac{1}{3}n$

(b) 220


(c) 20

7.6.2.17.**Answer.**


(a)

(b) $[-968, 972]$ **7.6.2.19.****Answer.**(a) $2, -1$


(b)

**7.6.2.21.****Answer.**(a) $0, 1, -3$


(b)

**7.6.2.23.****Answer.**(a) $0, 1, -1$


(b)

**7.6.2.25.****Answer.**(a) $-1, 1$

(b)

**7.6.2.27.****Answer.**(a) $-1, 1, \pm\sqrt{6}$

(b)

**7.6.2.29.****Answer.** $x(x+2)(x-3)$ **7.6.2.33.****Answer.** $x^2(x+4)(x-4)$ **7.6.2.35.****Answer.**(a) $P(-2) = 0$ **7.6.2.31.****Answer.** $x^3(x+2)(x-2)$ (b) $\frac{3 \pm \sqrt{13}}{2}$

7.6.2.37.

Answer. $-2 \pm i\sqrt{6}$

7.6.2.39.

Answer. $1 \pm \frac{\sqrt{6}}{3}i$

7.6.2.41.

Answer.

(a) -8 (b) -8

7.6.2.43.

Answer. $\frac{11}{10} - \frac{13}{10}i$



7.6.2.45.

Answer.

 $x^4 - 2x^3 + 14x^2 - 18x + 45$

7.6.2.47.

Answer.




7.6.2.49.

Answer.

(a) $V = \frac{\pi h^3}{4}$

(b) $2\pi \text{ cm}^3 \approx 6.28 \text{ cm}^3$; $16\pi \text{ cm}^3 \approx 50.27 \text{ cm}^3$


(c)



7.6.2.51.

Answer.

(a)




- (b) 338
 (c) Months 2 and 20
 (d) During month 6. The number of members eventually decreases to zero.

7.6.2.53.

Answer. All numbers except $-2, 0, 2$.

7.6.2.55.


Answer.

**7.6.2.57.**

Answer.

- (a) Horizontal asymptote $y = 0$;
 Vertical asymptote $x = 4$;
 y -intercept $(0, \frac{-1}{4})$


(b)

**7.6.2.59.**

Answer.

- (a) Horizontal asymptote $y = 1$;
 Vertical asymptote $x = -3$;
 x -intercept $(2, 0)$; y -intercept $(0, \frac{-2}{3})$


(b)

**7.6.2.61.**

Answer.

- (a) Horizontal asymptote $y = 3$;
 Vertical asymptote $x = \pm 2$;
 x -intercept $(0, 0)$; y -intercept $(0, 0)$


(b)



7.6.2.63.

Answer.


(a) $y = \frac{-5}{x+3} + 3$



7.6.2.65.

Answer.

(a) $y = \frac{2}{(x+1)^2} + 1$



7.6.2.67.

Answer.

(a) $t_1 = \frac{90}{v-2}$


(b) $t_2 = \frac{90}{v+2}$

(c)

7.6.2.69.

Answer.

(a)



7.6.2.71.

Answer. 299

7.6.2.73.


Answer. -2

7.6.2.77.

Answer. All a except -1 and 1

7.6.2.81.

Answer. $n = \frac{Ct}{C-V}$



(d) $\frac{90}{v-2} + \frac{90}{v+2} = 24$

(e) 8 mph

(b) $\frac{320}{x} = \frac{1}{2}x + 6$; \$20

7.6.2.75.

Answer. No solution

7.6.2.79.

Answer. 0

7.6.2.83.

Answer. $q = \frac{pr}{r-p}$

8 · Models and data

8.1 · Linear Regression

8.1.6 · Linear Regression (Homework 8.1)

8.1.6.1.

Answer.

a

x	50	125
y	9000	15,000

b $C = 5000 + 80x$

c $m = 80$ dollars/bike, so it costs the company \$80 per bike it manufactures.

8.1.6.3.

Answer.

a

g	12	5
d	312	130

b $d = 26g$

c $m = 26$ miles/gallon, so the Porche's fuel efficiency is 26 miles per gallon.

8.1.6.5.

Answer.

a


C	15	-5
F	59	23

b $F = 32 + \frac{9}{5}C$

c $m = \frac{9}{5}$, so an increase of 1°C is equivalent to an increase of $\frac{9}{5}^\circ\text{F}$.


8.1.6.7.

Answer.



8.1.6.9.

Answer.




8.1.6.11.

Answer.

a 12 seconds

b 39




d 11.6 seconds

e $y = 8.5 + 0.1x$

f 12.7 seconds; 10.18 seconds; The prediction for the 40-year-old is reasonable, but not the prediction for the 12-year-old.

8.1.6.13.

Answer.




b $y = 121 + 19.86t$

c 419

8.1.6.15.

Answer.




b $y = 64.2 - 1.63t$

c 58 births per 1000 women

d 32 births per 1000 women

8.1.6.17.

Answer.




b $y = 0.18t + 67.9$

c 74.9 years

d 79 years

8.1.6.19.

Answer.



b $y = 90.49t - 543.7$


c 90.49 dollars/year: Each additional year of education corresponds to an additional \$90.49 in weekly earnings.

d No: The degree or diploma attained is more significant than the number of years. So, for example, interpolation for the years of education between a bachelor's and master's degree may be inaccurate because earnings with just the bachelor's degree will not change until the master's degree is attained. And the years after the professional degree will not add significantly to earnings, so extrapolation is inappropriate.

8.1.6.21.

Answer.

a



b $y = 1.6 + 0.11t$

c 6.2 billion tons

8.1.6.23.**Answer.**


a 0.34 meters per year

b $y = 0.34x$ ($b = 0$ because the plant has zero size until it begins.)

c Over 1300 years

8.1.6.25.**Answer.**

a




b Yes

c $y = 1.29x - 1.62$

d The slope, 1.29 kg/sq cm, tells us that strength increases by 1.29 kg when the muscle cross-sectional area increases by 1 sq cm.

8.1.6.27.**Answer.**

a E

c $y = 1.33x$; There should be no loss in mass when no gas evaporates.

d 1333 mg

e Oxygen


8.1.6.29.**Answer.**

a 75°F

b The slope of -2 degrees/hour says that temperatures are dropping at a rate of 2° per hour.**8.1.6.31.****Answer.**

a 20 mph

b The slope of 10 mph/second says the car accelerates at a rate of 10 mph per second.

8.1.6.33.**Answer.** 2 min: 21°C ; 2 hr: 729°C ; The estimate at 2 minutes is reasonable; the estimate at 2 hours is not reasonable.**8.1.6.35.****Answer.** 128 lb.**8.1.6.37.****Answer.**a $y \approx -0.54x + 58.7$ 


b 31.7%

c 90 meters

d The regression line gives a negative probability, which is not reasonable.

8.1.6.39.**Answer.**

a



$$y \approx 22.8x + 198.5$$

b ≈ 540 watts

c 198.5 watts

d ≈ -8.7 newtons

e 3.5 watts

f about 0.018 or 1.8%

8.2 · Curve Fitting

8.2.5 · Curve-fitting (Homework 8.2)

8.2.5.1.

Answer. $a = -2, b = 3, c = -4$

8.2.5.3.

Answer. $a = 1, b = -4, c = 7$

8.2.5.5.

Answer. $a = 3, b = 1, c = -2$. The equation for the parabola is $y = 3x^2 + x - 2$


8.2.5.7.

Answer.

(a) $P = -0.16x^2 + 7.4x - 71$

(b) 14%. It predicts that 14% of the 25-year old population use marijuana on a regular basis.


(c)



8.2.5.9.

Answer.

(a) $C = 0.75t^2 - 1.85t + 56.2$



(b) 65.7 lb

8.2.5.11.

Answer. $D = \frac{1}{2}n^2 - \frac{3}{2}n$

8.2.5.13.

Answer.

(a) $y = a(x + 2)^2 + 6$

(b) 3

8.2.5.15.

Answer.

(a) $y = \frac{3}{4}x^2 - 3$

(b) $y = ax^2 - 3$ for any $a < 0$

8.2.5.17.

Answer. $y = -2(x - 30)^2 + 280$

8.2.5.19.

Answer. $y = x^2 - 9$

8.2.5.23.

Answer. $y = x^2 - 2x - 15$

8.2.5.21.

Answer. $y = -2x^2$

8.2.5.25.

Answer. $y = x^2 - 4x + 5$

8.2.5.27.**Answer.**

(a) $y = \frac{-1}{40}(x - 80)^2 + 164$

(b) 160.99 ft

8.2.5.29.**Answer.**

(a) Vertex: $(\frac{1991}{2}, 79)$; y -intercept: $(0, 297)$

(b) $y = 0.00022(x - 995.5)^2 + 79$

8.2.5.31.**Answer.**

(a) 8 m

(b) $y = \frac{x^2}{32}$


(c) 3.125 m

8.2.5.33.**Answer.**


(a) $h = 8.24t + 38.89$

(e) 135.7 m

(b) 162.5 m



(d) $h = -0.81t^2 + 21.2t$



(g) Quadratic: Gravity will slow the projectile, giving the graph a concave down shape.

8.2.5.35.**Answer.**

(a) $y = -0.587t^2 + 7.329t - 2.538$


(b) The predicted peak was in 2000, near the end of March. The model predicts 7 deaths for 2005.

8.2.5.37.**Answer.**


(a) $y = 0.0051t + 11.325$

(e) 7.4 hr

(b) 13.2 hr



(d) $y = -0.00016t^2 + 0.053t + 9.319$



(g) 9.8 hr (the same as the previous year); Neither model is appropriate.

Appendix G

GNU Free Documentation License

Version 1.3, 3 November 2008

Copyright © 2000, 2001, 2002, 2007, 2008 Free Software Foundation, Inc.
(<http://www.fsf.org/>)

Everyone is permitted to copy and distribute verbatim copies of this license document, but changing it is not allowed.

0. PREAMBLE. The purpose of this License is to make a manual, textbook, or other functional and useful document “free” in the sense of freedom: to assure everyone the effective freedom to copy and redistribute it, with or without modifying it, either commercially or noncommercially. Secondly, this License preserves for the author and publisher a way to get credit for their work, while not being considered responsible for modifications made by others.

This License is a kind of “copyleft”, which means that derivative works of the document must themselves be free in the same sense. It complements the GNU General Public License, which is a copyleft license designed for free software.

We have designed this License in order to use it for manuals for free software, because free software needs free documentation: a free program should come with manuals providing the same freedoms that the software does. But this License is not limited to software manuals; it can be used for any textual work, regardless of subject matter or whether it is published as a printed book. We recommend this License principally for works whose purpose is instruction or reference.

1. APPLICABILITY AND DEFINITIONS. This License applies to any manual or other work, in any medium, that contains a notice placed by the copyright holder saying it can be distributed under the terms of this License. Such a notice grants a world-wide, royalty-free license, unlimited in duration, to use that work under the conditions stated herein. The “Document”, below, refers to any such manual or work. Any member of the public is a licensee, and is addressed as “you”. You accept the license if you copy, modify or distribute the work in a way requiring permission under copyright law.

A “Modified Version” of the Document means any work containing the Document or a portion of it, either copied verbatim, or with modifications and/or translated into another language.

A “Secondary Section” is a named appendix or a front-matter section of the Document that deals exclusively with the relationship of the publishers or authors of the Document to the Document’s overall subject (or to related matters) and contains nothing that could fall directly within that overall subject. (Thus, if the Document is in part a textbook of mathematics, a Secondary Section may not explain any mathematics.) The relationship could be a matter of historical connection with the subject or with related matters, or of legal, commercial, philosophical, ethical or political position regarding them.

The “Invariant Sections” are certain Secondary Sections whose titles are designated, as being those of Invariant Sections, in the notice that says that the Document is released under this License. If a section does not fit the above definition of Secondary then it is not allowed to be designated as Invariant. The Document may contain zero Invariant Sections. If the Document does not identify any Invariant Sections then there are none.

The “Cover Texts” are certain short passages of text that are listed, as Front-Cover Texts or Back-Cover Texts, in the notice that says that the Document is released under this License. A Front-Cover Text may be at most 5 words, and a Back-Cover Text may be at most 25 words.

A “Transparent” copy of the Document means a machine-readable copy, represented in a format whose specification is available to the general public, that is suitable for revising the document straightforwardly with generic text editors or (for images composed of pixels) generic paint programs or (for drawings) some widely available drawing editor, and that is suitable for input to text formatters or for automatic translation to a variety of formats suitable for input to text formatters. A copy made in an otherwise Transparent file format whose markup, or absence of markup, has been arranged to thwart or discourage subsequent modification by readers is not Transparent. An image format is not Transparent if used for any substantial amount of text. A copy that is not “Transparent” is called “Opaque”.

Examples of suitable formats for Transparent copies include plain ASCII without markup, Texinfo input format, LaTeX input format, SGML or XML using a publicly available DTD, and standard-conforming simple HTML, PostScript or PDF designed for human modification. Examples of transparent image formats include PNG, XCF and JPG. Opaque formats include proprietary formats that can be read and edited only by proprietary word processors, SGML or XML for which the DTD and/or processing tools are not generally available, and the machine-generated HTML, PostScript or PDF produced by some word processors for output purposes only.

The “Title Page” means, for a printed book, the title page itself, plus such following pages as are needed to hold, legibly, the material this License requires to appear in the title page. For works in formats which do not have any title page as such, “Title Page” means the text near the most prominent appearance of the work’s title, preceding the beginning of the body of the text.

The “publisher” means any person or entity that distributes copies of the Document to the public.

A section “Entitled XYZ” means a named subunit of the Document whose title either is precisely XYZ or contains XYZ in parentheses following text that translates XYZ in another language. (Here XYZ stands for a specific section name mentioned below, such as “Acknowledgements”, “Dedications”, “Endorsements”, or “History”.) To “Preserve the Title” of such a section when you modify the Document means that it remains a section “Entitled XYZ” according to this definition.

The Document may include Warranty Disclaimers next to the notice which states that this License applies to the Document. These Warranty Disclaimers

are considered to be included by reference in this License, but only as regards disclaiming warranties: any other implication that these Warranty Disclaimers may have is void and has no effect on the meaning of this License.

2. VERBATIM COPYING. You may copy and distribute the Document in any medium, either commercially or noncommercially, provided that this License, the copyright notices, and the license notice saying this License applies to the Document are reproduced in all copies, and that you add no other conditions whatsoever to those of this License. You may not use technical measures to obstruct or control the reading or further copying of the copies you make or distribute. However, you may accept compensation in exchange for copies. If you distribute a large enough number of copies you must also follow the conditions in section 3.

You may also lend copies, under the same conditions stated above, and you may publicly display copies.

3. COPYING IN QUANTITY. If you publish printed copies (or copies in media that commonly have printed covers) of the Document, numbering more than 100, and the Document's license notice requires Cover Texts, you must enclose the copies in covers that carry, clearly and legibly, all these Cover Texts: Front-Cover Texts on the front cover, and Back-Cover Texts on the back cover. Both covers must also clearly and legibly identify you as the publisher of these copies. The front cover must present the full title with all words of the title equally prominent and visible. You may add other material on the covers in addition. Copying with changes limited to the covers, as long as they preserve the title of the Document and satisfy these conditions, can be treated as verbatim copying in other respects.

If the required texts for either cover are too voluminous to fit legibly, you should put the first ones listed (as many as fit reasonably) on the actual cover, and continue the rest onto adjacent pages.

If you publish or distribute Opaque copies of the Document numbering more than 100, you must either include a machine-readable Transparent copy along with each Opaque copy, or state in or with each Opaque copy a computer-network location from which the general network-using public has access to download using public-standard network protocols a complete Transparent copy of the Document, free of added material. If you use the latter option, you must take reasonably prudent steps, when you begin distribution of Opaque copies in quantity, to ensure that this Transparent copy will remain thus accessible at the stated location until at least one year after the last time you distribute an Opaque copy (directly or through your agents or retailers) of that edition to the public.

It is requested, but not required, that you contact the authors of the Document well before redistributing any large number of copies, to give them a chance to provide you with an updated version of the Document.

4. MODIFICATIONS. You may copy and distribute a Modified Version of the Document under the conditions of sections 2 and 3 above, provided that you release the Modified Version under precisely this License, with the Modified Version filling the role of the Document, thus licensing distribution and modification of the Modified Version to whoever possesses a copy of it. In addition, you must do these things in the Modified Version:

- A. Use in the Title Page (and on the covers, if any) a title distinct from that of the Document, and from those of previous versions (which should, if

there were any, be listed in the History section of the Document). You may use the same title as a previous version if the original publisher of that version gives permission.

- B. List on the Title Page, as authors, one or more persons or entities responsible for authorship of the modifications in the Modified Version, together with at least five of the principal authors of the Document (all of its principal authors, if it has fewer than five), unless they release you from this requirement.
- C. State on the Title page the name of the publisher of the Modified Version, as the publisher.
- D. Preserve all the copyright notices of the Document.
- E. Add an appropriate copyright notice for your modifications adjacent to the other copyright notices.
- F. Include, immediately after the copyright notices, a license notice giving the public permission to use the Modified Version under the terms of this License, in the form shown in the Addendum below.
- G. Preserve in that license notice the full lists of Invariant Sections and required Cover Texts given in the Document's license notice.
- H. Include an unaltered copy of this License.
- I. Preserve the section Entitled "History", Preserve its Title, and add to it an item stating at least the title, year, new authors, and publisher of the Modified Version as given on the Title Page. If there is no section Entitled "History" in the Document, create one stating the title, year, authors, and publisher of the Document as given on its Title Page, then add an item describing the Modified Version as stated in the previous sentence.
- J. Preserve the network location, if any, given in the Document for public access to a Transparent copy of the Document, and likewise the network locations given in the Document for previous versions it was based on. These may be placed in the "History" section. You may omit a network location for a work that was published at least four years before the Document itself, or if the original publisher of the version it refers to gives permission.
- K. For any section Entitled "Acknowledgements" or "Dedications", Preserve the Title of the section, and preserve in the section all the substance and tone of each of the contributor acknowledgements and/or dedications given therein.
- L. Preserve all the Invariant Sections of the Document, unaltered in their text and in their titles. Section numbers or the equivalent are not considered part of the section titles.
- M. Delete any section Entitled "Endorsements". Such a section may not be included in the Modified Version.
- N. Do not retitle any existing section to be Entitled "Endorsements" or to conflict in title with any Invariant Section.
- O. Preserve any Warranty Disclaimers.

If the Modified Version includes new front-matter sections or appendices that qualify as Secondary Sections and contain no material copied from the Document, you may at your option designate some or all of these sections as invariant. To do this, add their titles to the list of Invariant Sections in the Modified Version's license notice. These titles must be distinct from any other section titles.

You may add a section Entitled "Endorsements", provided it contains nothing but endorsements of your Modified Version by various parties — for example, statements of peer review or that the text has been approved by an organization as the authoritative definition of a standard.

You may add a passage of up to five words as a Front-Cover Text, and a passage of up to 25 words as a Back-Cover Text, to the end of the list of Cover Texts in the Modified Version. Only one passage of Front-Cover Text and one of Back-Cover Text may be added by (or through arrangements made by) any one entity. If the Document already includes a cover text for the same cover, previously added by you or by arrangement made by the same entity you are acting on behalf of, you may not add another; but you may replace the old one, on explicit permission from the previous publisher that added the old one.

The author(s) and publisher(s) of the Document do not by this License give permission to use their names for publicity for or to assert or imply endorsement of any Modified Version.

5. COMBINING DOCUMENTS. You may combine the Document with other documents released under this License, under the terms defined in section 4 above for modified versions, provided that you include in the combination all of the Invariant Sections of all of the original documents, unmodified, and list them all as Invariant Sections of your combined work in its license notice, and that you preserve all their Warranty Disclaimers.

The combined work need only contain one copy of this License, and multiple identical Invariant Sections may be replaced with a single copy. If there are multiple Invariant Sections with the same name but different contents, make the title of each such section unique by adding at the end of it, in parentheses, the name of the original author or publisher of that section if known, or else a unique number. Make the same adjustment to the section titles in the list of Invariant Sections in the license notice of the combined work.

In the combination, you must combine any sections Entitled "History" in the various original documents, forming one section Entitled "History"; likewise combine any sections Entitled "Acknowledgements", and any sections Entitled "Dedications". You must delete all sections Entitled "Endorsements".

6. COLLECTIONS OF DOCUMENTS. You may make a collection consisting of the Document and other documents released under this License, and replace the individual copies of this License in the various documents with a single copy that is included in the collection, provided that you follow the rules of this License for verbatim copying of each of the documents in all other respects.

You may extract a single document from such a collection, and distribute it individually under this License, provided you insert a copy of this License into the extracted document, and follow this License in all other respects regarding verbatim copying of that document.

7. AGGREGATION WITH INDEPENDENT WORKS. A compilation of the Document or its derivatives with other separate and independent

documents or works, in or on a volume of a storage or distribution medium, is called an “aggregate” if the copyright resulting from the compilation is not used to limit the legal rights of the compilation’s users beyond what the individual works permit. When the Document is included in an aggregate, this License does not apply to the other works in the aggregate which are not themselves derivative works of the Document.

If the Cover Text requirement of section 3 is applicable to these copies of the Document, then if the Document is less than one half of the entire aggregate, the Document’s Cover Texts may be placed on covers that bracket the Document within the aggregate, or the electronic equivalent of covers if the Document is in electronic form. Otherwise they must appear on printed covers that bracket the whole aggregate.

8. TRANSLATION. Translation is considered a kind of modification, so you may distribute translations of the Document under the terms of section 4. Replacing Invariant Sections with translations requires special permission from their copyright holders, but you may include translations of some or all Invariant Sections in addition to the original versions of these Invariant Sections. You may include a translation of this License, and all the license notices in the Document, and any Warranty Disclaimers, provided that you also include the original English version of this License and the original versions of those notices and disclaimers. In case of a disagreement between the translation and the original version of this License or a notice or disclaimer, the original version will prevail.

If a section in the Document is Entitled “Acknowledgements”, “Dedications”, or “History”, the requirement (section 4) to Preserve its Title (section 1) will typically require changing the actual title.

9. TERMINATION. You may not copy, modify, sublicense, or distribute the Document except as expressly provided under this License. Any attempt otherwise to copy, modify, sublicense, or distribute it is void, and will automatically terminate your rights under this License.

However, if you cease all violation of this License, then your license from a particular copyright holder is reinstated (a) provisionally, unless and until the copyright holder explicitly and finally terminates your license, and (b) permanently, if the copyright holder fails to notify you of the violation by some reasonable means prior to 60 days after the cessation.

Moreover, your license from a particular copyright holder is reinstated permanently if the copyright holder notifies you of the violation by some reasonable means, this is the first time you have received notice of violation of this License (for any work) from that copyright holder, and you cure the violation prior to 30 days after your receipt of the notice.

Termination of your rights under this section does not terminate the licenses of parties who have received copies or rights from you under this License. If your rights have been terminated and not permanently reinstated, receipt of a copy of some or all of the same material does not give you any rights to use it.

10. FUTURE REVISIONS OF THIS LICENSE. The Free Software Foundation may publish new, revised versions of the GNU Free Documentation License from time to time. Such new versions will be similar in spirit to the present version, but may differ in detail to address new problems or concerns. See <http://www.gnu.org/copyleft/>.

Each version of the License is given a distinguishing version number. If the Document specifies that a particular numbered version of this License “or

any later version” applies to it, you have the option of following the terms and conditions either of that specified version or of any later version that has been published (not as a draft) by the Free Software Foundation. If the Document does not specify a version number of this License, you may choose any version ever published (not as a draft) by the Free Software Foundation. If the Document specifies that a proxy can decide which future versions of this License can be used, that proxy’s public statement of acceptance of a version permanently authorizes you to choose that version for the Document.

11. RELICENSING. “Massive Multiauthor Collaboration Site” (or “MMC Site”) means any World Wide Web server that publishes copyrightable works and also provides prominent facilities for anybody to edit those works. A public wiki that anybody can edit is an example of such a server. A “Massive Multiauthor Collaboration” (or “MMC”) contained in the site means any set of copyrightable works thus published on the MMC site.

“CC-BY-SA” means the Creative Commons Attribution-Share Alike 3.0 license published by Creative Commons Corporation, a not-for-profit corporation with a principal place of business in San Francisco, California, as well as future copyleft versions of that license published by that same organization.

“Incorporate” means to publish or republish a Document, in whole or in part, as part of another Document.

An MMC is “eligible for relicensing” if it is licensed under this License, and if all works that were first published under this License somewhere other than this MMC, and subsequently incorporated in whole or in part into the MMC, (1) had no cover texts or invariant sections, and (2) were thus incorporated prior to November 1, 2008.

The operator of an MMC Site may republish an MMC contained in the site under CC-BY-SA on the same site at any time before August 1, 2009, provided the MMC is eligible for relicensing.

ADDENDUM: How to use this License for your documents. To use this License in a document you have written, include a copy of the License in the document and put the following copyright and license notices just after the title page:

```

    Copyright (C) YEAR YOUR NAME.
    Permission is granted to copy, distribute and/or modify this
document
    under the terms of the GNU Free Documentation License, Version
1.3
    or any later version published by the Free Software Foundation;
    with no Invariant Sections, no Front-Cover Texts, and no
Back-Cover Texts.
    A copy of the license is included in the section entitled "GNU
    Free Documentation License".

```

If you have Invariant Sections, Front-Cover Texts and Back-Cover Texts, replace the “with... Texts.” line with this:

```

    with the Invariant Sections being LIST THEIR TITLES, with the
    Front-Cover Texts being LIST, and with the Back-Cover Texts
being LIST.

```

If you have Invariant Sections without Cover Texts, or some other combination of the three, merge those two alternatives to suit the situation.

If your document contains nontrivial examples of program code, we recommend releasing these examples in parallel under your choice of free software license, such as the GNU General Public License, to permit their use in free software.

Bibliography

- [1] Ahrens, C. Donald. *Essentials of Meteorology*. Belmont, CA: Wadsworth, 1998.
- [2] Alexander, R. M. *The Human Machine*. Natural History Museum Publications, 1992.
- [3] Altenburg, Edgar. *Genetics*. New York: Holt Rinehart and Winston, 1956.
- [4] Atkins, P. W. and Beran, J. A. *General Chemistry*. Scientific American Books, 1989.
- [5] Baddeley, Alan D. *Essentials of Human Memory*. Psychology Press Ltd., 1999.
- [6] Belovsky, G. E. "Diet Optimization in a Generalist Herbivore: The Moose." *Theoretical Population Biology*, 14 (1978), 105–134.
- [7] Belovsky, G. E. "Generalist Herbivore Foraging and Its Role in Competitive Interactions." *American Zoologist*, 26 (1986), 51–69.
- [8] Bender, Edward. *An Introduction to Mathematical Modeling*. Mineola, NY: Dover Publications, 1978.
- [9] Berg, Richard and Stork, David. *The Physics of Sound*. Englewood Cliffs, NJ: Prentice-Hall, 1982.
- [10] Boleman, Jay. *Physics, a Window on Our World*. Englewood Cliffs, NJ: Prentice Hall, 1982.
- [11] Bolton, William. *Patterns in Physics*. McGraw Hill, 1974.
- [12] Brandt, John C. and Maran, Stephen P. *New Horizons in Astronomy*. San Francisco: W. H. Freeman, 1972.
- [13] Briggs, David and Courtney, Frank. *Agriculture and Environment*. Singapore: Longman Scientific and Technical, 1985.
- [14] Burton, R. F. *Biology by Numbers*. Cambridge: Cambridge University Press, 1998.
- [15] Carr, Bernard, and Giddings, Steven. "Quantum Black Holes." *Scientific American*, May 2005.
- [16] Chapman J. L. and Reiss M. J. *Ecology: Principles and Applications*. Cambridge: Cambridge University Press, 1992.
- [17] Davis, Kimmet, and Autry. *Physical Education: Theory and Practice*. Macmillan Education Australia PTY Ltd., 1986.
- [18] Deffeyes, Kenneth. *Hubbert's Peak*. Princeton: Princeton University Press, 2001.

- [19] *The Earth and Its Place in the Universe*. Open University, Milton Keynes, 1998.
- [20] Fowler, Murray and Miller, Eric. *Zoo and Wild Animal Medicine*. Saunders, 1999.
- [21] Gillner, Thomas C. *Modern Ship Design*. U.S. Naval Institute Press, 1972.
- [22] *The Guinness Book of Records*, 1998. Guinness Publishing Ltd., 1997.
- [23] Hayward, Geoff. *Applied Ecology*. Thomas Nelson and Sons, Ltd., 1992.
- [24] Holme, David J., and Peck, Hazel. *Analytical Biochemistry*. Longman Scientific and Technical, 1993.
- [25] Hunt, J. A. and Sykes, A. *Chemistry*. Longman Group Ltd., 1984.
- [26] Hutson, A. H. *The Pocket Guide to Mammals of Britain and Europe*. Dragon's World Ltd., 1995.
- [27] Ingham, Neil. *Astrophysics*. International Thomson Publishing, 1997.
- [28] Karttunen, H. et al. *Fundamental Astronomy*. Springer-Verlag, 1987.
- [29] Krebs J. R. and Davies N. B. *An Introduction to Behavioural Ecology*. Oxford: Blackwell Scientific Publications, 1993.
- [30] Leopold, Luna B., Wolman, M. Gordon, and Miller, John P. *Fluvial Processes in Geomorphology*. New York: Dover Publications, Inc., 1992.
- [31] Mannering, Fred, and Kilareski, Walter. *Principles of Highway Engineering and Traffic Analysis*. New York: John Wiley and Sons, Inc., 1998.
- [32] Meadows, Donella, Randers, Jorgen, and Meadows, Dennis. *Limits to Growth, the 30-Year Update*. White River Junction, VT: Chelsea Green Publishing Co., 2004.
- [33] Mee, Frederick. *Sound*. Heinemann Educational Books, 1967.
- [34] Oglesby, Clarkson and Hicks, R. Gary. *Highway Engineering*. New York: John Wiley and Sons, Inc., 1982.
- [35] Oliver, C. P. "The Effect of Varying the Duration of X-Ray Treatment upon the Frequency of Mutation," *Science*, 71 (1930), 44–46.
- [36] Perrins, C. M. *British Tits*. Collins, London, 1979.
- [37] Perrins, C. M. and Moss, D. "Reproductive Rates in the Great Tit." *Journal of Animal Ecology*, 44 (1975), 695–706.
- [38] Plummer, David. *Biochemistry, the Chemistry of Life*. London: McGraw-Hill Book Co., 1989.
- [39] Pope, Jean A. *Medical Physics*. Heinemann Educational, 1989.
- [40] P. E. di Prampero et al. *Journal of Applied Physiology*, 37 (1974), 1–5.
- [41] Pratt, Paul W. *Principles and Practices of Veterinary Technology*. Mosby, 1998.
- [42] Pugh, L. G. C. E. "Relation of Oxygen Intake and Speed in Competition Cycling and Comparative Observations on the Bicycle Ergometer." *Journal of Physiology*, 241 (1974), 795–808
- [43] Scarf, Philip. "An Empirical Basis for Naismith's Rule." *Mathematics Today*, vol. 34 no. 5 (1998), 149–151
- [44] Schad, Jerry. *Afoot and Afield in Los Angeles County*. Berkeley: Wilder-

- ness Press, 1991.
- [45] Schmidt-Nielsen, Knut. *How Animals Work*. Cambridge: Cambridge University Press, 1972.
 - [46] Schmidt-Nielsen, Knut. *Scaling: Why Is Animal Size so Important?* Cambridge: Cambridge University Press, 1984.
 - [47] Sears, Francis. *Mechanics, Heat, and Sound*. Sternberg, Robert J. In Search of the Human Mind. Harcourt Brace College Publishing, 1995.
 - [48] Storch, Hammon, and Bunch. *Ship Production*. Cornell Maritime Press, 1988.
 - [49] Strickberger, Monroe W. *Genetics*. Macmillan, 1976.
 - [50] Underwood, Benton J. "Forgetting." *Scientific American*, vol. 210, no. 3, 91–99.
 - [51] Weisberg, Joseph and Parish, Howard. *Introductory Oceanography*. McGraw-Hill, 1974.
 - [52] Wilkinson, G. S. "Reciprocal Food Sharing in the Vampire Bat". *Nature*, 308 (1984), 181–184.
 - [53] Wood, Alexander. *The Physics of Music*. Chapman and Hall, 1975.
 - [54] Wright, Paul and Paquette, Radnor. *Highway Engineering*. New York: John Wiley and Sons, Inc., 1987.

Index

- absolute value, 170
- additive inverse, 972
- algebraic equation, 3
- algebraic expression, 866
- algebraic fraction, 918
- allometric equation, 339
- allometric equations, 290
- allometry, 339
- amortize, 479
- amplitude, 282
- annuity, 477
- aphelion, 386
- ascending powers, 698
- average, 870
- axis of symmetry, 633

- base, 443, 946
- bell-shaped curve, 594
- binomial, 903
- biological half-life, 803
- body mass index, 811
- build, 924
- building factor, 924
- Bézier curves, 695

- carrying capacity, 594
- Cartesian, 880
- center, 951
- change of base, 596
- circle, 951
- closed interval, 862
- common denominator, 923
- common logarithms, 445
- complex conjugate, 739
- complex number, 624, 737
- complex numbers, 341, 624, 640, 737
- complex plane, 742
- compound inequality, 862
- compounded continuously, 560
- concavity, 216

- conjugate, 937
- conjugate pair, 916
- conjugates, 916
- constant of variation, 292
- continuous compounding, 551
- coordinate system, 880
- counting numbers, 970
- cube root, 852
- cubic, 173, 715

- decay factor, 400, 402, 490
- decibel scale, 577
- decreasing graph, 7
- degree, 904
- dependent, 28
- descending powers, 698
- difference of two squares, 915
- direct variation, 291
- directly proportional, 291
- discriminant, 640, 643, 678
- distance formula, 950
- domain, 250
- doubling time, 473

- e, 552
- effective half-life, 803
- elimination, 832, 888
- empirical model, 219
- equilibrium price, 783, 869
- equivalent equations, 859
- error tolerance, 241
- evaluating, 866
- evaluating the function, 33
- experience curve, 380
- exponential equation, 421
- exponential function, 491
- exponential functions, 420
- exponential growth, 403, 490
- extraction of roots, 154
- extraneous, 539, 938
- extraneous solution, 778

- extrapolation, 814
- factored form, 603
- factoring, 905
- fatigue index, 387
- formula, 861
- function, 28, 250
- functions, 14
- fundamental theorem of algebra, 741
- future value, 477, 478, 480, 493, 505
- general form, 9
- golden rectangle, 785
- graph, 3
- graph of an equation, 880
- growth factor, 394, 402, 490
- half-closed, 863
- half-life, 475
- half-open, 863
- horizontal asymptote, 174, 752, 756
- horizontal line test, 515
- hypotenuse, 945
- imaginary axis, 742
- imaginary part, 624, 737
- imaginary unit, 623, 735
- increasing graphs, 7
- independent, 28
- index of the radical, 333
- inequality, 861
- inequality symbol, 861
- infinite intervals, 863
- inflection point, 594, 715
- initial value, 394, 402
- input, 28
- integers, 970
- intercept, 633
- intercepts, 11
- intercepts of the graph, 11
- interpolation, 814
- interval, 862
- inverse function, 508
- inverse function notation, 510
- inverse square law, 320
- inverse variation, 298
- inversely proportional, 299
- irrational numbers, 335, 971
- isosceles, 946
- laws of exponents, 321, 895
- LCD, 923
- lead coefficient, 698
- least squares regression line, 815
- like fractions, 922
- line of best, 811
- linear combination, 890
- linear combinations, 890
- linear equations, 9
- linear growth, 403
- linear models, 3
- linear regression, 813
- linear system, 887
- log-log graph paper, 575
- logarithm, 443
- logarithmic equation, 537
- logarithmic function, 533
- logarithmic scale, 572
- logistic function, 594
- lowest common denominator, 923
- mean, 870
- mechanistic models, 220
- midline, 282
- midpoint, 950
- monomial, 903
- multiplicative inverse, 973
- multiplicity, 607
- natural base, 551, 552, 561
- natural exponential function, 552, 561
- natural log function, 554, 561
- natural logarithm, 553
- natural numbers, 970
- negative exponents, 316
- nonstrict, 862
- n th roots, 333
- one-to-one, 515
- open interval, 863
- opposite, 972
- output, 28
- parabola, 173, 633
- parameters, 110
- percent decrease, 869
- percent increase, 398, 869
- perfect-square trinomials, 915
- perihelion, 386
- period, 282
- periodic function, 279
- periods, 279
- pH scale, 576
- piecewise, 177
- point-slope form, 114
- polynomial, 903
- polynomial function, 698

- power function, 318
- power functions, 289, 354, 420
- present value, 479, 480, 493, 505
- proportional, 291
- pure imaginary, 624, 737

- quadratic formula, 620
- quadratic function, 601, 609
- quadratic in, 608
- quadratic in form, 607
- quartic, 717

- radical, 333
- radical equation, 340
- radical notation, 357
- radicand, 333
- radius, 951
- range, 250
- rate of change, 88
- ratio method, 472
- rational expression, 918
- rational function, 749, 759, 790
- rational functions, 174
- rational number, 970
- rationalizing the denominator, 936
- real axis, 742
- real numbers, 970
- real part, 624, 737
- reciprocal, 973
- rectangular, 880
- regression line, 813
- repeater bar, 971
- Richter, 579
- right triangle, 944

- scales, 374
- scaling, 297

- scaling exponent, 297
- scatterplot, 811
- scientific notation, 853
- sigmoid function, 594
- similar, 947
- sinusoidal, 282
- slope, 85, 86, 108
- slope-intercept form, 108
- solution, 880, 887
- standard form, 603, 951
- strict inequalities, 862
- subscripts, 90
- substitution, 887
- supergrowth, 381
- surface area, 949

- table of values, 3
- transformations, 190
- translations, 190
- triangle inequality, 946
- trinomial, 903
- two-parameter, 110

- union, 863

- varies directly, 295
- varies inversely, 298, 299
- varies jointly, 375
- vertex, 633
- vertex form, 660
- vertical asymptote, 174, 750, 754, 759, 790
- volume, 948

- weighted average, 870
- whole numbers, 970

Colophon

This book was authored in PreTeXt.